

AN OVERVIEW ON PHYTOCHEMISTRY AND ETHNOMEDICINAL POTENTIAL OF *ACANTHUS ILICIFOLIUS* OF WEST BENGAL REGION

Biswajit Pradhan¹, Keshamma E², Kamal Kant Patra^{3*}, Veermani Kumar⁴

¹Ph.D. Scholar, Department of Botany, YBN University, Ranchi, Jharkhand, India

²Associate Professor (Ph.D. Co-Supervisor), Department of Biochemistry,
Maharani Cluster University, Palace Road, Bengaluru, Karnataka, India

³Associate Professor, Department of Botany, YBN University, Ranchi, Jharkhand, India

⁴Ph.D. Supervisor, Assistant Professor, Department of Botany, YBN University,
Ranchi, Jharkhand, India

***Correspondence Author**

Dr. Kamal Kant Patra

Associate Professor,
Department of Botany,
YBN University, Ranchi,
Jharkhand- 834 010

India

Email ID: kamalnalbot@rediffmail.com

Abstract

Mangroves are one among the most productive and biologically important ecosystem on this planet, providing vital ecosystem goods and services. *Acanthus ilicifolius* generally known as Sea holy and holy mangrove. The genus *Acanthus*, belonging to the family Acanthaceae, is most common in both east and west coast of India including Andaman Islands and Nicobar Islands. Genus *Acanthus* comprises of over 30 species, However, in India only *A. ilicifolius* is known to occur in all mangrove habitats. *A. ilicifolius* is common in landward edges of mangroves just above the high tide mark, and also occurs in inner mangroves. Out of all the species, *A. ilicifolius* has a long history of use as a folklore and traditional remedy for treating various diseases in Ayurveda as well as in Thai medicine and Traditional Chinese Medicine. Furthermore, all the parts of *A. ilicifolius* plant have been used to cure a number of ailments in various traditional systems of medicine including Ayurveda. Thus, in the present review we aimed to describe and delineate on the geographical distribution, phytochemistry and ethnomedicinal properties of *A. ilicifolius*.

Keywords: Mangroves, *Acanthus ilicifolius*, Phytochemistry, Ayurveda, Ethnomedicinal properties

Introduction

Acanthus ilicifolius generally known as Sea holy and holy mangrove. Mangroves are considered as one of the most specialized ecological assemblages of halophytic plants acting as a transient zone between land and ocean. They comprise of taxonomically diverse shrubs and trees, distributed along tropical and sub-tropical environments having specific habitats such as shores, estuaries, tidal creeks, backwaters, lagoons, marshes, mudflats and even at upstream points where water remains saline.¹ Mangrove forests are unique functional ecosystems having much social, economic and biological importance. They are among one of the most productive ecosystems of the world as they provide important ecosystem supplies and services to human society as well as coastal and marine systems.^{2,3} These habitats interact with a wide array of aquatic or terrestrial flora and fauna, enabling their growth and establishment. Considering their value for the environment and coastal communities, mangrove conservation should become a priority and efforts must be invested to find new and successful methods for conserving mangrove ecosystems.⁴

The genus *Acanthus*, belonging to the family Acanthaceae, comprises of over 30 species, mostly shrubs or perennial herbs, native of tropics and subtropics, many of which are found from India to southern China, tropical Australia and the Western Pacific islands, throughout Southeast Asia.⁵ In India, it is common in both east and west coast including Andaman Islands and Nicobar Islands. The name '*Acanthus*' is derived from a Greek word meaning 'spiny' and the genus is also known as 'Bear's Breeches' due to the spiny nature of the plant. The genus comprises of mangrove species viz. *Acanthus ebracteatus* Vahl., *A. ilicifolius* L., *A. volubilis* Wall., *A. latisepalus*, *A. montanus* and *A. xiamenensis* that are known to sustain in the most hostile atmosphere making this genus a unique taxon among all true mangrove genera that represent both terrestrial and true mangrove species.

A. xiamenensis is endemic to China and all the other species are common in Indo-West Pacific region. However, in India only *A. ilicifolius* is known to occur in all mangrove habitats except Lakshadweep, whereas *A. ebracteatus* and *A. volubilis* have very restricted distribution. *A. ilicifolius* is common in landward edges of mangroves just above the high tide mark, and also occurs in inner mangroves. In ancient times, *Acanthus* leaves were employed as a decorative motif in architecture of the Mediterranean countries. In the late 1960s, the founders of The Arts Society chose the *Acanthus* leaf as the Society's emblem. Out of all the species, *A. ilicifolius* has a long history of use as a folklore and traditional remedy for treating various diseases in Ayurveda as well as in Thai medicine and Traditional Chinese Medicine.^{6,7} Hence, this genus represents medicinally as well as economically important plants.⁸ With this scenario, in the present review we aimed to describe and delineate on the geographical distribution, phytochemistry and ethnomedicinal properties of *A. ilicifolius*.

Taxonomy of *A. ilicifolius*

Class: Dicotyledones

Subclass: Gamopetalae

Series: Bicarpellatae

Order: Personales

Family: Acanthaceae

Genus: *Acanthus*Species: *ilicifolius***Environment and Geographical Distribution of *A. ilicifolius***

A. ilicifolius is regular in estuaries all through Asian tropics from India to Polynesia and northern Australia. Asia and Australia have the best assortment and conveyance of mangrove species known to mankind. In India, it is reported from the east (the largest area of mangrove forest, the Sunderbans), the West seashore, Andamans and a North eastern state, Meghalaya as depicted in Figure 1.⁹ It typically develops on waterway banks or tidal channel sides or low swampy zones in mangrove woodlands and regions.¹⁰



Figure 1: Showing geographical distribution of *A. ilicifolius*

Morphology of *A. ilicifolius*

A. ilicifolius is a shrub grows up to 2 meter tall in robust mangrove areas, Aerial roots-stilt roots; Leaves-simple, opposite, lanceolate, narrowed at base, serrate margins armed with spines; spines longer in flowering season; petiole short, 5 to 6 mm in length, slightly winged with two sharp spurs at the base; color dark green when fresh, yellowish-brown on drying. Odor-indistinct, taste bitter. Flowers-sessile, 4 cm long, spike inflorescence, terminal, corolla light blue or violet; Fruit-capsule, ovoid-oblong, up to 3 cm long, compressed, apiculate, brown, shining (Figure 2).^{7,11}



Figure 2: Showing *A. ilicifolius* plant

Microscopic Characteristics of *A. ilicifolius*

Leaf can be characterized by important features like presence of salt glands with a unicellular stalk and multicellular head in adaxial side; median vein with numerous irregular collateral vascular bundles; the lateral vein bundles somewhat near to median bundle with continuous layer of sclerenchyma sheath; the two layered palisade tissue being confined on abaxial side only; glandular trichomes and deeply sunken stomata intermittently seen in the abaxial region.¹¹ Stem shows single layered thick epidermis; outer cortex with collenchyma cells and inner cortex consisting of parenchyma with air cavities; solitary sclereids embedded in peripheral phloem cells; pith made up of small parenchymatous cells that contain numerous calcium oxalate crystals and water-storing large parenchymatous cells. The diagnostic features of powdered drug include glandular trichomes with 2 to 10 celled salt secreting head either sessile or with 1 to 2-celled stalk; lower epidermis with plenty of diacytic stomata; transversely cut fragments of lamina exhibiting upper epidermis with sunken pits embedded with glandular trichomes; groups of long pericyclic fibres with thick pitted wall and narrow lumen; and pitted lignified parenchymatous cells.¹²

Phytochemistry of *A. ilicifolius*

Studies on phytochemical investigation of the genus *Acanthus* indicated that phytochemical study has been extensively done on the true mangrove species; *A. ilicifolius*, beside this work no phytochemical isolation has been performed on this genus.¹³ Through several phytochemical studies, *A. ilicifolius* has been explored for the presence of various classes of bioactive compounds namely alkaloids, benzoxazinoids, lignans, flavonoids, triterpenoids and steroids.¹⁴ The phytochemicals reported in various parts of *A. ilicifolius* are described as below:

Phytochemistry of leaf

The alkaloids reported in leaves of *A. ilicifolius* are as follows:

- Acanthicifoline, a 1-methyl-1,2,3,4-tetrahydro-5-methoxy-2,7-naphthyridin-3-one¹⁵
- Trigonellin¹⁶
- 2-benzoxazolinone^{17,18}
- Acanthiline A (pyrido[1,2-a] indole); 4-hydroxybenzoxazole-2-one¹⁹
- Indole alkaloids like 1H-indole-3-carboxylic acid, 1H-indole-3-carboxaldehyde and 1H-indole-3-acetic acid²⁰
- Betaine²¹

The Benzoxazinoids reported in leaves of *A. ilicifolius* are as follows:

- Benzoxazin-3-one, benzoxazinoidglucosides²²
- 4-benzoxazin-3(4H)-one, megastigmane 2-benzoxazolinone, 5,5'-bis-benzoxazoline-2,2'-dione¹²

The lignans reported in leaves of *A. ilicifolius* are as follows:

- A coumaric acid derivative called acancifoliuside²³
- Acteoside²⁴
- Isoacteoside²⁵
- Acanthaminoside²⁶
- (+)-syringaresinol-O-β-D-glucopyranoside, (+)-lyoniresinol-3α-O-β-glucopyranoside²⁷
- (-)-lyoniresinol are isolated from the methanolic extract of leaf; (+)-lyoniresinol 3α-O-β-D-galactopyranosyl-(1→6)-β-D-glucopyranoside²⁸
- (+)-lyoniresinol-2α-O-β-D-galactopyranosyl-3α-O-β-D-glucopyranoside have been reported from aerial parts²⁹
- (Z)-4-coumaric acid glycosides like (Z)-4-coumaric aci-4-O-β-D-glucopyranoside and (Z)-4-coumaric acid-4-O-β-D-apiofuranosyl-(1'→2')-O-β-D-glucopyranoside⁵
- Phenylethyl-O-β-D-glucopyranosyl-(1→2)-β-D-glucopyranoside, phenylethyl-O-β-D-glucopyranoside, benzyl-O-β-D-glucopyranoside and vanillic acid²¹
- 411-epoxymegastigmane glucoside and megastigmane glucosides [(6S, 9S)-roseoside]) have also been reported from *A. ilicifolius* growing in China⁵
- The phenylethanoids reported in the plant are blepharin, acteoside, isoverbascoside³⁰
- Martynoside and crenatoside³¹

The flavonoids reported in leaves of *A. ilicifolius* are as follows:

- Quercetin, quercetin-3-O-β-D-glucopyranoside³²
- Apigenin-7-O-β-D-glucuronide, vitexin³³
- Luteolin 7-O-β-D-glucuronide, apigenin-7-O-β-D-glucuronide, methylapigenin-7-O-β-D-glucuronate³³
- Catechin/epicatechin, galatechin/epicatechin and amphicin/epigalin³

The aliphatic glycosides reported in leaves of *A. ilicifolius* are as follows:

- Ilifolioside B³³
- Ilicifolioside C³⁴

The triterpenoids reported in leaves of *A. ilicifolius* are as follows:

- Pentacyclic triterpenes, β -amyrin, α -amyrin³⁵
- Lupeol, oleanolic acid and ursolic acid¹⁶

The steroids reported in leaves of *A. ilicifolius* are as follows:

- β -Sitosterol³⁵
- Cholesterol, campesterol, stigmasterol³⁶
- Stigmast-7-en-3- β -ol, octacosyl alcohol, stigmasteryl β -D-glucopyranoside³⁷
- Daucosterol and 3-O- β -D-glucopyranosyl-stigmasterol³⁰

The hydrocarbons reported in leaves of *A. ilicifolius* are as follows:

- 24-Methylene cycloartenol, cycloartenol, 19-Norlanost-5-en-3-ol, 24-methylenelanost-9-en-3 β -ol, 4-methyl-cholesta-7-ene-3 β -ol, phytol, octadecanoate, tetracosanol and octacosanol^{36,38}

The fatty acid derivatives reported in leaves of *A. ilicifolius* are as follows:

- Palmitic acid and octadecanoic acid³⁶

Other phytochemicals reported in leaves of *A. ilicifolius* are as follows:

- Amino acids like glutamic and aspartic acids and organic acids like malic, citric, oxalic glycolic and succinic acids³⁹
- Uridine and uracil⁴⁰

Phytochemistry of roots

The major phytochemicals reported in roots of *A. ilicifolius* are as follows:

- Triterpenoids⁴¹
- 2-benzoxazolinone (acanthosides A-D)⁴²

Phytochemistry of pods

The major phytochemicals reported in pods of *A. ilicifolius* are as follows:

- 1,4-Benzoxazinone⁵

Phytochemistry of stems

The major phytochemicals reported in stems of *A. ilicifolius* are as follows:

- Homologous series of 15 saturated odd and even fatty acids⁴³
- Lignan and cyclolignan glycoside^{29,44}
- Phenylethanoid glycoside (ilicifolioside A), an aliphatic alcohol glycoside-ilicifolioside B²³

- Lignin glucosides, (+)-lyoniresinol-3 α -[2-(3,5-dimethoxy-4-hydroxy)-benzoyl]-O- β -glucopyranoside and dihydroxymethyl-bis (3,5-dimethoxy-4-hydroxyphenyl) tetrahydrofuran-9(or 9')-O- β -glucopyranoside; and benzoxazinoidglucosides^{12,15}

Ethnomedicinal Properties of *A. ilicifolius*

The pharmacological study has been extensively performed on *A. ilicifolius* among the species of this genus. This study revealed that *A. ilicifolius* has significant bioactivities like, antioxidant and cytotoxic activity,⁴⁵ antinociceptive activity,⁴⁶ anti-inflammatory activity,⁴⁷ anti-osteoporotic activity,³⁵ hepatoprotective activity,⁴⁷ chemopreventive activity; leishmanicidal, tumour reducing and anticancer activities;^{24,25,47,48} antileishmanial, antiulcer, antimicrobial, and osteoblastic, activities.⁴⁷ Moreover, Firdaus et al. found in their study that flower extract of the plant also has good cytotoxic effects;⁴⁵ Smitha et al. showed the anticancer activity of ethyl acetate extract of on two cell lines PA-1 and MCF-7.⁴⁹ The major ethnopharmaceutical activities of *A. ilicifolius* described in details herein:

Anti-inflammatory

The methanolic fraction of leaf extract of *A. ilicifolius* is reported to produce significant inhibition of rat paw oedema, when administered both prior to and after carrageenan administration, in a manner similar to BW755C, a synthetic cyclooxygenase (COX) and lipoxygenase (LOX) inhibitor. The extract is shown to decrease protein exudation and leukocyte migration in the peritoneal fluid, thereby indicating its effectiveness towards inhibiting peritoneal inflammation and also significant inhibition of COX (1 and 2) and 5-LOX activity. Moreover, pre-incubation of the extract is shown to inhibit the production of proinflammatory cytokines (TNF- α and IL-6) in lipopolysaccharide (LPS)-stimulated peripheral blood mononuclear cells (PBMCs). It is also reported to possess significant free radical (DPPH, ABTS, superoxide and hydroxyl radical) scavenging activity.⁵⁰

Antioxidant and Cytotoxic

The methanol extract of *A. ilicifolius* contains phenolic substances grouped as cancer prevention agent compounds.⁵¹ The limit of antiradical proficiency of *A. ilicifolius* flowers can be as medium. The extract was not in unadulterated structure; be that as it may, it may very well be arranged as a decent and potential antioxidant agent. The LC₅₀ of extracts of *A. ilicifolius* or pure compounds on salt water shrimp or cell line under 100 μ g/mL is ordered as a potential cytotoxic and dangerous substance.⁵² The ethanol leaves extract of *A. ilicifolius* was seen as cytotoxic towards lung fibroblast (L-929) cells in 72 h MTT test and the focus required for half-cell passing was 18 μ g/mL,²⁵ in the interim the methanol extract of *A. ilicifolius* was cytotoxic to Hela and κ B cell line.⁵³ Wostmann and Liebezeit detailed that this mangrove contained cancer prevention agent substances.⁵⁴ The most noteworthy cancer prevention agent and cytotoxic activity were found on methanol extract. Methanol has been known progressively powerful to break up dynamic mixes in cells. Henceforth, it was simpler to infiltrate the cell film to extract the intracellular fixings from plant materials. Tiwari et al. expressed that few dynamic mixes will be gotten if methanol utilized as dissolvable in the extraction strategy for example anthocyanins,

saponins, tannins, flavones, and polyphenols.⁵⁵ These mixes have known as free radical scavenger, responsive species quencher, hydrogen donor, cancer prevention agent chemicals activator, detoxification inducer, ordinary normal cell differentiation promoter and expansion cell inhibitor, and apoptosis inducer.⁵⁶⁻⁶⁰

The cytotoxicity of methanol extract of *A. ilicifolius* can be related to the antioxidant activity and synergism impact of multi-part in separate. Triterpenoid saponin demonstrated its cytotoxicity in HeLa cells through both mitochondrial brokenness and ER stress cell passing pathways, while saponin stifled tumor obtrusive and relocation by hindering MMP-2 and MMP-9 activation.⁶¹ Imai et al. established that flavonoid adequately smothered the multiplication of a human colon carcinoma cell line, COLO 201, through apoptosis acceptance while phenolic indicated anticancer action on disease colon cell by capturing the cell cycle.⁶² All in all, the methanol concentrate of *A. ilicifolius* flower is potential as antioxidant (cancer prevention agent) substances and cytotoxic compounds.⁴⁵

Furthermore, Firdaus et al., analyzed the cytotoxic of the flower extracts of *A. ilicifolius* on the brine shrimp lethality. The results exhibited methanol extract has lowest LC₅₀ value (22 µg/ml) while water extract showed the highest value at 10 µg/ml among the extracts.⁴⁵ Dey et al. have tested the cytotoxic activity of a methanolic crude extract of the plant where it has found that the extract exhibited a significant cytotoxic activity with the LC₅₀ value at 60 µg/ml and LC₉₀ value at 120 µg/ml.⁶³

Antimicrobial

Antimicrobial activity of alcohol, butanol and chloroform extract (10 mg/ml) of both leaves and roots of the *A. ilicifolius* exhibited significant activity against all microbes, *B. subtilis*, *P. vulgaris*, *P. aeruginosa*, *S. aureus*, *C. albicans*, *A. niger*, and *A. fumigatus*. The experiment was followed by agar cup plate method that unveils the leaves were more active than that of roots where ampicillin used as a standard for bacteria and clotrimazole for fungi. Chloroform extract of leaf showed the highest zone of inhibition that was measured ~26 mm against the fungus *C. albicans*. But butanol extract of leaf extract displayed lowest ~08 mm against the bacterium *S. aureus*.⁶⁴

Govindasamy and Arulpriya studied antimicrobial activity of *A. ilicifolius* against seven skin infection causing microbes, methicillin resistance *S. aureus* (MRSA), *L. plantarum*, *S. pyogenes*, *S. epidermis*, *C. albicans*, *P. aeruginosa*, and *T. rubrum*. In this test four extract of the leaf has been examined where chloroform extract showed the highest activity against all microbes (except *S. epidermis* and *L. plantarum*).⁴⁹

Antiviral

The ethanol extracts (50%) of the leaf, flower and root of *A. ilicifolius* have been shown to possess in vitro antiviral activity against Tobacco Mosaic Virus.^{65,66} Blepharin, acteoside, isoverbascoside, daucosterol and 3-O-β-D-glucopyranosyl-stigmasterol isolated from the leaf when investigated for anti-influenza virus activities by measuring the neuraminidase activity of

influenza virus, the phenylethanoids (blepharin, acteoside, isoverbascoside) exhibited significant antiviral activities.³⁰

Anti-leishmanial

2-Benzoxazolinone (BOA) isolated from the leaf (40 µg/ml) of *A. ilicifolius* is shown to have anti-leishmanial activity against *Leishmania donovani* when compared with pentamine in in vitro experiments. BOA (i.p.) in the doses ranging from 0.25 to 1 g/kg is shown to be safe in mice.^{65,67}

Antinociceptive

The methanolic extract of *A. ilicifolius* is reported to exhibit antinociceptive effects, due to its flavonoids and tannins, and is mediated by inhibition of lipooxygenase and/or cyclooxygenase in peripheral tissues induced by acetic acid, thereby reducing PGE2 synthesis and interfering with the mechanism of transduction in primary afferent nociceptor.^{68,69}

Anticancer

Smitha et al. studied the anticancer activity of ethyl acetate extract of leaf and root extract of *A. ilicifolius* on two cell lines, MCF-7 and PA-1. *A. ilicifolius* is more effective on PA-1 and the result has recommended that at 50 µg/ml ratio is adequate to inhibit the cancer cells. Moreover, the result has also unveiled that the extract is slightly cytotoxic to both of the cell lines.⁷⁰ Benzoxazolinone-type alkaloids, acanthosides A-D yielded from the root of *A. ilicifolius* are reported to possess significant cytotoxicity against the HepG2, HeLa, and A-549 cancer cell lines with IC₅₀ 7.8 to 26.6 µM, due to the presence of substituted benzoyl moiety in their structures.⁷¹

Hepatoprotective

The alcoholic extract (250 and 500 mg/kg, p.o.) of *A. ilicifolius* has shown significant hepatoprotective activity in rats in carbon tetrachloride-induced hepatotoxicity model, that was comparable with curcumin (100 mg/kg, p.o.), as evaluated from the serum and tissue activity of marker enzymes.²⁴

Antidiabetic

A single oral dose of 200 and 400 mg/kg of ethanolic extract of root is shown to significantly decrease blood glucose levels in alloxan-induced diabetic rats after 5 h and 3 h respectively in an acute study and on first day in a sub-acute study. It is reported to improve regeneration of β-cells of pancreas.⁷²

Osteoblastic

Osteoblasts are the bone-framing cells of the skeleton; they incorporate, direct and regulate the mineralization of the extra-cell grid of the bone. MC3T3-E1 cells, an osteoblast-like cell line, have been accounted for to hold their ability to separate into osteoblasts, and may give valuable data on the impacts of phytochemicals on the separation of osteoblasts.⁷³ Acancifoliuside, a coumaric acid derivative isolated from the methanolic leaf extract of *A. ilicifolius*, when tested for its effects on the functions of osteoblastic MC3T3-E1 cells, significantly increased the growth and differentiation of osteoblasts, indicating that *A. ilicifolius* leaf may help to prevent osteoporosis.³⁵

Anti-ulcer

The anti-ulcer activity of methanolic leaf extract of the *A. ilicifolius* (at 100 to 200 mg/kg) produces significant inhibition of gastric lesions induced by pylorus ligation and ethanol-induced gastric ulcers in Wistar albino rats. The extract is shown to cause significant reduction in the gastric volume, free acidity and ulcer index as compared to control, probably due to anti-secretory action.⁶⁶

Conclusion

The world's tropical and subtropical regions are home to a large dispersion of *Acanthus* plants. The ethnomedicinal applications *Acanthus* in health care management have received remarkable recognition as a result of systematic pharmacological studies on the species. A true mangrove plant, *A. ilicifolius* is found throughout the world's tropical and subtropical regions. *A. ilicifolius* botanical research offers scientific data on morphological and anatomical traits.

The phytochemical studies revealed different bioactive compounds from the various parts of *A. ilicifolius*. Numerous pharmacological studies done using different types of extracts of *A. ilicifolius* revealed various pharmacological properties such as antimicrobial, anti-inflammatory, antiviral, antidiabetic, antioxidant, cytotoxic and anticancer activities, antinociceptive, hepatoprotective, leishmanicidal and osteoblastic activity which encouraged the use of *A. ilicifolius* in ethnomedicine. Thus, more research on *A. ilicifolius* is advised in order to cover the advantages of this mangrove plant and uncover hidden areas to support clinical applications for improved health.

References

1. Qasim SZ. Glimpses of the Indian Ocean. Universities Press; 1998:123-9.
2. Bouillon S, Dahdouh-Guebas F, Rao AV, Koedam N, Dehairs F. Sources of organic carbon in mangrove sediments: variability and possible ecological implications. *Hydrobiologia*. 2003; 495:33-9.
3. FAO, 2007. The World's Mangroves 1980-2005. FAO Forestry Paper No. 153. Rome, Forest Resources Division, FAO.77.
4. Bosold AL. Challenging the “man” in mangroves: the missing role of women in mangrove conservation.2012.
5. Wu J, Zhang S, Xiao Q, Li Q, Huang J, Long L, Huang L. Phenylethanoid and aliphatic alcohol glycosides from *Acanthus ilicifolius*. *Phytochemistry*. 2003;63(4):491-5.
6. Wostmann R, Liebezeit G. Chemical composition of the mangrove holly *Acanthus ilicifolius* (Acanthaceae)—review and additional data. *Senckenbergiana maritima*. 2008;38(1):31-7.
7. Saranya A, Ramanathan T, Kesavanarayanan KS, Adam A. Traditional medicinal uses, chemical constituents and biological activities of a mangrove plant, *Acanthus ilicifolius* Linn.: a brief review. *Am. Eurasian J. Agric. Environ. Sci*. 2015;15(2):243-50.

8. Bhattacharyya B, Johri BM. Flowering plants: taxonomy and phylogeny. 1998.
9. Weblink1: Available from: <https://www.discoverlife.org/mp/20m?kind=Acanthus+ilicifolius> (Accessed 10 May 2024).
10. Wang Y, Zhu H, Tam NF. Polyphenols, tannins and antioxidant activities of eight true mangrove plant species in South China. *Plant and soil*. 2014; 374:549-63.
11. Surya S, Hari N. Anatomical, morphological, palynological, phytochemical and molecular profiling of medicinal mangrove *Acanthus ilicifolius* L. (Acanthaceae). *Star Internat J.*, 2018; 6(4): 41-2.
12. Gupta AK, Tandon N, Sharma M. Quality Standards of Indian Medicinal Plants. 1st Edn., New Delhi; Indian Council of Medical Research, 2008.
13. Bora R, Adhikari PP, Das AK, Raaman N, Sharma GD. Ethnomedicinal, phytochemical, and pharmacological aspects of genus *Acanthus*. *Int J Pharm Pharm Sci*. 2017;9(12):8.
14. Patel R, Patel N, Patel K, Patel M, Patel K, Verma P, Shah M. *Acanthus ilicifolius*: A true mangrove with biomedical potential. *WJPPS*. 2020;9(11):471-89.
15. Tiwari KP, PK M, Masood M. Acanthicifoline-a new alkaloid from *Acanthus ilicifolius*. *Pol J Chem*, 1980; 54: 857-8.
16. Tan D, Jiang C, Tao Y. Chemical constituents of *Acanthus ilicifolius*. *Chemistry of Natural Compounds*. 2016;52(5):951-2.
17. Minocha PK, Tiwari KP. A triterpenoidal saponin from roots of *Acanthus illicifolius*. *Phytochemistry*. 1981;20(1):135-7.
18. Murty MS, Solimabi S, Kamat SY. Isolation of 2-benzoxazolinone from *Acanthus ilicifolius*. *Indian J Pharm Sci.*, 1984; 46: 218-9
19. Long S, Wei X, Chen Y, Zheng G, Li, X, Peng X, Li S. Preparation and application of *Acanthus ilicifolius* alkaloid A and its derivative with anti-inflammatory and analgesic activities. *Faming Zhuanli Shenqing Gongkai Shuomingshu*, 2009; CN101397279: A 20090401.
20. Liu Q, Lin W. Study on chemical constituents of *Acanthus ilicifolius* L. of (sic) mangrove plant from southsea (sic) China. *Journal of Harb in University of Commerce (Natural Sciences Edition)*. 2008; 6:648-51.
21. Huo CH, Wang B, Liang H, Zhao YY, Lin WH. Study on chemical constituents of mangrove *Acanthus ilicifolius*. *China journal of Chinese materia medica*. 2006; 31(24): 2052-4.
22. Kokpol U, Chittawong V, Miles DH. Chemical constituents of the roots of *Acanthus illicifolius*. *Journal of Natural Products*. 1986;49(2):355-6.
23. Kanchanapoom T, Kasai R, Yamasaki K. Flavonoid glycosides from *Acanthus ilicifolius* L. (Natural Medicine Note). *Natural medicines*. 2002;56(3):122.
24. Babu BH, Shylesh BS, Padikkala J. Antioxidant and hepatoprotective effect of *Acanthus ilicifolius*. *Fitoterapia*. 2001;72(3):272-7.

25. Babu BH, Shylesh BS, Padikkala J. Tumour reducing and anticarcinogenic activity of *Acanthus ilicifolius* in mice. *Journal of Ethnopharmacology*. 2002;79(1):27-33.
26. Bich DH, Trung DQ, Chuong BX, Dong NT, Dam DT, Hien PV, Lo VN, Mai PD, Man PK, Nhu DT, Tap N. The medicinal plants and animals of Vietnam, Hanoi Science and Technology Publisher, Hanoi. Vol. II. 2004:635-6.
27. Chi VV. The Dictionary of Vietnamese Medicinal Plants. 1st Edn., Hanoi; Medicine Publisher: 1999.
28. Fang H, Tong W, Shi LM, Blair R, Perkins R, Branham W, Hass BS, Xie Q, Dial SL, Moland CL, Sheehan DM. Structure– activity relationships for a large diverse set of natural, synthetic, and environmental estrogens. *Chemical research in toxicology*. 2001;14(3):280-94.
29. Wu J, Zhang S, Li Q, Huang J, Xiao Z, Long L. Two new cyclolignan glycosides from *Acanthus ilicifolius*. *Zeitschrift für Naturforschung B*. 2004;59(3):341-4.
30. Chen YP, Tan DP, Zeng Q, Wang Y, Yan QX, Zeng LJ. Chemical constituents from leaves of *Acanthus ilicifolius* and their anti-influenza virus activities. *Journal of Chinese Medicinal Materials*. 2015; 38(3): 527-30.
31. Zhang M, Ren X, Yue S, Zhao Q, Shao C, Wang C. Simultaneous quantification of four phenylethanoid glycosides in rat plasma by UPLC-MS/MS and its application to a pharmacokinetic study of *Acanthus ilicifolius* herb. *Molecules*. 2019;24(17):3117.
32. Tan D, Jiang C, Tao Y. Chemical constituents of *Acanthus ilicifolius*. *Chemistry of Natural Compounds*. 2016;52(5):951-2.
33. Wu J, Zhang S, Xiao Q, Li Q, Huang J, Long L, Huang L. Megastigmane and flavone glycosides from *Acanthus ilicifolius*. *Die Pharmazie-An International Journal of Pharmaceutical Sciences*. 2003;58(5):363-4.
34. Ragavan P, Saxena A, Mohan PM, Jayaraj RS, Ravichandran K. Taxonomy and distribution of species of the genus *Acanthus* (Acanthaceae) in mangroves of the Andaman and Nicobar Islands, India. *Biodiversitas Journal of Biological Diversity*. 2015;16(2).
35. Van Kiem P, Quang TH, Huong TT, Nhung LT, Cuong NX, Van Minh C, Choi EM, Kim YH. Chemical constituents of *Acanthus ilicifolius* L. and effect on osteoblastic MC3T3E1 cells. *Archives of Pharmacal Research*. 2008; 31:823-9.
36. Zhong LJ, Huang MY, Zhang JG, Li GW, Zhang YH. Study on the chemical constituents from *Acanthus ilicifolius* Linn. var. *xiamenensis*. *Chin J Mar Drugs*. 2012;31(06):23-8.
37. Kokpol U, Chittawong V, Miles DH. Chemical constituents of the roots of *Acanthus illicifolius*. *Journal of Natural Products*. 1986;49(2):355-6.
38. Hai F, Tan XI, Li GQ. Steroids and terpenoids from the mangrove *Acanthus ilicifolius*. *Tianran Chanwu Yanjiu Yu Kaifa*, 2010; 22(4): 597-9.

39. Wu J, Zhang S, Huang J, Xiao Q, Li Q, Long L, Huang L. New aliphatic alcohol and (Z)-4-coumaric acid glycosides from *Acanthus ilicifolius*. Chemical and pharmaceutical bulletin. 2003;51(10):1201-3.
40. Huo CH, Zhao YY, Liang H, Lin WH. Studies on chemical constituents in herbs of *Acanthus ilicifolius*. China journal of Chinese materia medica. 2005 May;30(10):763-5.
41. Minocha PK, Tiwari KP. A triterpenoidal saponin from roots of *Acanthus ilicifolius*. Phytochemistry. 1981;20(1):135-7.
42. Zhao D, Xie L, Yu L, An N, Na W, Chen F, Li Y, Tan Y, Zhang X. New 2-Benzoxazolinone derivatives with cytotoxic activities from the roots of *Acanthus ilicifolius*. Chemical and Pharmaceutical Bulletin. 2015;63(12):1087-90.
43. Goswami S, Chatterjee B, Mallik M. Proof-of-presence of unusually naturally occurring homologous series of fifteen saturated odd and even fatty acids in *Acanthus ilicifolius* L. (Acanthaceae). Journal of the Indian Chemical Society. 2004;81(8):696-706.
44. Kanchanapoom T, Kamel MS, Kasai R, Yamasaki K, Picheansoonthon C, Hiraga Y. Lignan glucosides from *Acanthus ilicifolius*. Phytochemistry. 2001;56(4):369-72.
45. Firdaus M, Prihanto AA, Nurdiani R. Antioxidant and cytotoxic activity of *Acanthus ilicifolius* flower. Asian Pacific Journal of Tropical Biomedicine. 2013;3(1):17-21.
46. Islam MA, Saifuzzaman M, Ahmed F, Rahman MM, Sultana NA, Naher K. Antinociceptive activity of methanolic extract of *Acanthus ilicifolius* Linn leaves. Turk J Pharm Sci. 2012;9(1):51-60.
47. Ganesh S, Vennila JJ. Screening for antimicrobial activity in *Acanthus ilicifolius*. Arch Appl Sci Res. 2010; 2:311–31.
48. De Carvalho PB, Ferreira EI. Leishmaniasis phytotherapy. Nature's leadership against an ancient disease. Fitoterapia. 2001;72(6):599-618.
49. Govindasamy C, Arulpriya M. Antimicrobial activity of *Acanthus ilicifolius*: Skin infection pathogens. Asian Pacific Journal of Tropical Disease. 2013;3(3):180-3.
50. Kumar KM, Gorain B, Roy DK, Samanta SK, Pal M, Biswas P, Roy A, Adhikari D, Karmakar S, Sen T. Anti-inflammatory activity of *Acanthus ilicifolius*. Journal of Ethnopharmacology. 2008;120(1):7-12.
51. Spavieri J, Allmendinger A, Kaiser M, Casey R, Hingley-Wilson S, Lalvani A, Guiry MD, Blunden G, Tasdemir D. Antimycobacterial, antiprotozoal and cytotoxic potential of twenty-one brown algae (phaeophyceae) from British and Irish waters. Phytotherapy Research. 2010;24(11):1724-9.
52. Ara J, Sultana V, Ehteshamul-Haque S, Qasim R, Ahmad VU. Cytotoxic activity of marine macro-algae on *Artemia salina* (Brine shrimp). Phytotherapy Research: An International Journal Devoted to Pharmacological and Toxicological Evaluation of Natural Product Derivatives. 1999;13(4):304-7.
53. Khajure PV, Rathod JL. Potential anticancer activity of *Acanthus ilicifolius* extracted from the mangroves forest of Karwar, West coast of India. World J Technol. 2011; 1:1-6.

54. Wostmann R, Liebezeit G. Chemical composition of the mangrove holly *Acanthus ilicifolius* (Acanthaceae)—review and additional data. *Senckenbergiana maritima*. 2008;38(1):31-7.
55. Tiwari P, Kumar B, Kaur M, Kaur G, Kaur H. Phytochemical screening and extraction: a review. *Internationale pharmaceutica scientia*. 2011;1(1):98-106.
56. Breton F, Cérantola S, Gall EA. Distribution and radical scavenging activity of phenols in *Ascophyllum nodosum* (Phaeophyceae). *Journal of Experimental Marine Biology and Ecology*. 2011;399(2):167-72.
57. Nielsen MF, Ingold KU. Kinetic solvent effects on proton and hydrogen atom transfers from phenols. Similarities and differences. *Journal of the American Chemical Society*. 2006;128(4):1172-82.
58. Fernández-Pachón MS, Berna G, Otaolauruchi E, Troncoso AM, Martín F, García-Parrilla MC. Changes in antioxidant endogenous enzymes (activity and gene expression levels) after repeated red wine intake. *Journal of agricultural and food chemistry*. 2009;57(15):6578-83.
59. Hsu YW, Tsai CF, Chen WK, Huang CF, Yen CC. A subacute toxicity evaluation of green tea (*Camellia sinensis*) extract in mice. *Food and Chemical Toxicology*. 2011;49(10):2624-30.
60. Chahar MK, Sharma N, Dobhal MP, Joshi YC. Flavonoids: A versatile source of anticancer drugs. *Pharmacognosy reviews*. 2011;5(9):1.
61. Man S, Gao W, Zhang Y, Huang L, Liu C. Chemical study and medical application of saponins as anti-cancer agents. *Fitoterapia*. 2010;81(7):703-14.
62. Imai M, Kikuchi H, Denda T, Ohyama K, Hirobe C, Toyoda H. Cytotoxic effects of flavonoids against a human colon cancer derived cell line, COLO 201: a potential natural anti-cancer substance. *Cancer Letters*. 2009;276(1):74-80.
63. Dey A, Raihan SM, Sariful Islam HM, Hamiduzzaman M, Monjur-Al-Hossain ASM. Phytochemical screening and the evaluation of the antioxidant, cytotoxic and antimicrobial properties of *Acanthus ilicifolius* (Family: Acanthaceae). *Int Res J Pharm* 2012; 3:153-6.
64. Bose S, Bose A. Antimicrobial activity of *Acanthus ilicifolius* (L.). *Indian journal of pharmaceutical sciences*. 2008;70(6):821.
65. Gupta AK, Tandon N. *Reviews on Indian Medicinal Plants*. 1st Edn., New Delhi; Indian Council of Medical Research: 2004.
66. Nizamuddin BS, Danamma B, Chitta S, Mohd D, Abdul M. Evaluation of antiulcer activity in the methanol extract of *Acanthus ilicifolius* leaves in experimental rats. *Int J Pharm Ind Res*. 2011; 1:57-62.
67. Kapil A, Sharma S, Wahidulla S. Leishmanicidal activity of 2-benzoxazolinone from *Acanthus illicifolius* in vitro. *Planta Medica*. 1994;60(02):187-8.

68. Collier HO, Dinneen LC, Johnson CA, Schneider C. The abdominal constriction response and its suppression by analgesic drugs in the mouse. *British journal of pharmacology and chemotherapy*. 1968;32(2):295.
69. Deraedt R, Jouquey S, Delevallée F, Flahaut M. Release of prostaglandins E and F in an algogenic reaction and its inhibition. *European journal of pharmacology*. 1980;61(1):17-24.
70. Smitha RB, Madhusoodanan PV, Prakashkumar R. Anticancer activity of *Acanthus ilicifolius* Linn. from chettuva mangroves, Kerala, India. *Int J Bioassays* 2014; 3:3452-5.
71. Zhao D, Xie L, Yu L, An N, Na W, Chen F, Li Y, Tan Y, Zhang X. New 2-Benzoxazolinone derivatives with cytotoxic activities from the roots of *Acanthus ilicifolius*. *Chemical and Pharmaceutical Bulletin*. 2015;63(12):1087-90.
72. Venkataiah G, Ahmed MI, Reddy DS, Rejeena M. Anti-diabetic activity of *Acanthus ilicifolius* root extract in alloxan induced diabetic rats. *Indo American Journal of Pharmaceutical Research*. 2013;3(11):9007-12.
73. Kodama HA, Amagai Y, Sudo H, Kasai S, Yamamoto S. Establishment of a clonal osteogenic cell line from newborn mouse calvaria. *Japanese Journal of Oral Biology*. 1981;23(4):899-901.