

ACCESSIBLE QUANTITATIVE TECHNIQUES FOR MEASURING FURANTHRIL IN BOTH PURE FORM AND IN PHARMACEUTICAL FORMULATIONS

Ajit Kumar Varma^{1*}, Rajat Srivastava², Neha Sirvastava², Gyanendra Singh²,
Nishant Singh Katiyar²

¹School of Pharmacy, Lingaya's Vidyapeeth, Faridabad, Haryana, 121002, India

²Faculty of Pharmaceutical Sciences, Rama University, Mandhana, Kanpur
(U.P.) India-209217

*Corresponding author: Ajit Kumar Varma, School of Pharmacy, Lingaya's Vidyapeeth,
Faridabad, Haryana, 121002, India

Abstract

Aims: Design of technical methods for the determination of Furanthril in its pure and pharmaceutical dosage form using spectral methods.

Study design: Planned and executed to estimate Furanthril by using Visible spectrophotometric in pure and pharmaceutical dosage form.

Methodology: Furanthril, the commercially known drug Lazix, which is important in the treatment of heart diseases and high blood pressure. This study was carried out using JASCO V – 630 double-beam computerized UV-Visible spectrophotometer with 1 cm matched cell, and HANA pH meter was used for reported pH readings.

Results: The reaction between Furanthril and bromo-phenol blue, xylenol orange, and chromazorol S. The decreasing in the intensity of the resulted colored complex was measured using bromo-phenol blue, xylenol orange, While the increasing of the color intensity was measured in the third method. These three method were based on charge transfer reaction. The limits of Beer's law for the first 0.4-32 μ g. mL⁻¹, second method 1-32 and the third method were 0.8-32 depending on the level of concentration, while the values of the molar absorption coefficient 1.4 \times 10⁴, 2.1 \times 10⁴ and 1.57 \times 10⁴ l.mol⁻¹.cm⁻¹ for the first, second and third method respectively. Sandel's significance also was calculated for these three methods, 0.0157 μ g.cm⁻² for the first method, 0.0236 μ g.cm⁻² for the second method, while the third method was 0.0210 μ g.cm⁻². The method has been successfully applied for the determination of Furanthril in its pure form and in some of its pharmaceutical preparations

Conclusion:The proposed method was validated in terms of linearity, range, Accuracy, precision, Specificity, Robustness. Method was successfully applied to the estimation of Furanthril, in its pure pharmaceutical dosage form.

Keywords:Furanthril, Xylenol Orange, Bromo-Phenol Blue, Chromazorol S, Pharmaceutical Preparations.

1. INTRODUCTION

Furanthril (FSD) is known commercially as Lasix, and chemically as 4-chloro-en-furfuryl-5-sulfamoylantranilic acid or 5-(aminesulfonyl)-4-chloro-2[(2-furanylmethyl) amino] benzoic acid. FSD is used to treat high blood pressure as well as a diuretic where it is used to treat edema associated with heart failure, cirrhosis, and kidney disease.[1-5].





Figure 1: Chemical structure of Furanthril


FSD is an important drug for human health, as it is used to treat the most important organs of the body, starting with the heart, kidneys and liver. Therefore, researchers have dealt with this drug in many studies, which have dealt with its solubility, uses and methods of estimation it, mostly of these method used various techniques, and among of these techniques, a spectrophotometric and chromatographic methods have been used for determination of FSD depending on: charge transfer complex method[6], mass spectroscopy [7], infra-red spectroscopy [8], Also, first order derivative spectroscopy method and absorbance ratio (Q-Absorbance) method were used.[9], other spectrophotometric method was using principal component regression[11], Validated RP-HPLC Method was used for estimation Furanthril in tablet[12], Some researchers depending on diazotized method to assay FSD[13], liquid chromatography were reported for estimation FSD in plasma and urine samples[14,15], as well as polarographic method[16]. Other method was based on using schiff's bases to estimate FSD spectrophotometrically [17].also, liquid-liquid extraction and high-performance liquid chromatography were used for estimation FSD[18,19]. Spectrophotometric methods adopted different reactions for the determination of FSD in pharmaceutical dosage forms [20-23], finally, reverse-phase high-performance liquid chromatography[24],and flow injection with HPLC[25], as well as HPLC method were used in the estimation of FSD [26-28]

In this paper, three an organic dyes have been used for estimating FSD, chromazurol S, xylenol orange and bromophenol blue, FSD was bleaching both of xylenol orange and bromophenol blue, where, chromazurol S was differed from them and increase the absorbance intensity as increasing FSD concentration, depending on this principle, FSD was assayed in three methods, method A bleaching bromophenol blue with increasing FSD amount at pH 3.63, as well as method B with xylenol orange at pH 4.72, While method C based on charge transfer reaction between chromazurol S and FSD at pH 4.77 with increasing FSD amount. So that, the organic dye used in method A is bromophenol blue which is chemically and

traditionally named as 3',3'',5',5''-tetrabromophenolsulfonphthalein, albutest respectively was prepared by the slowly addition of an excess bromine to a hot phenolsulfonphthalein in glacial acetic acid solution.[29], Xylenol orange is an organic reagent used as indicator for metal titration, Xylenol orange is the traditionally name of 2,2',2'',2'''-{(1,1-Dioxo-2-benzoxathiole-3,3 (1H)-diyl) bis [(6-hydroxy-5-methyl-3,1-phenylene) methylenenitrilo]}tetra acetic acid.[30]. Trisodium 5-[(E)-(3-carboxy-5-methyl-4-oxocyclohexa-2,5-dien-1-ylidene)(2,6-dichloro-3-sulfonatophenyl)methyl]-3-methyl-2-oxidobenzoate is the chemical name for chromazurol S, this reagent was widely used for determination of cations as well as medications in direct and indirect ways[31].



Bromophenol blue



Chromazurol

2. Experimental

Apparatus:

The final spectrum of FSD was measured and drawn using JASCO V – 630 double-beam computerized UV-Visible spectrophotometer, for all spectrophotometric measurements, 1 cm matched cell was used, and HANA pH meter used for reported pH readings.

Analytical chemicals

All chemicals used were of the purest analytical grade.

Table 1. Preparation of chemical materials

Chemical materials	Manufactured company	Weight, g	Solvent in final volum 100 mL	Concentration
FSD	SDI, Iraq	0.01	Ethanol[32]	100 µg/ ml
Bromophenol blue	Hopkin and williams	0.6699	Distilled water	0.01 M
Xylenol orange	Fluka	0.7585	Distilled water	0.01 M
Chromazurol S	Fluka	0.0605	Distilled water	0.001 M

Preparation of Furanthril from tablets

Three different brands of pharmaceutical preparations were used for Furanthril, where 10 tablets (each tablet contains 40 mg) were ground into a very fine powder, then weighed precisely about 0.01 g of the powder, this quantity was then dissolved in methanol, filtered and completed the volume of filtration with methanol mixed with warm distilled water at a ratio of 1:1 in a 100ml volumetric flask.

3. Discuss the experimental results:

100 µg of FSD in a final volume of 25 mL was used to study the experimental optimal conditions

Study of optimum conditions:

In this research paper, the optimal conditions suitable for the formation of the colored complex of Furanthril were studied and selected.

Selected the optimum medium of the reaction:

In order to choose the most appropriate type of acids and in the optimum quantity among sulfuric, acetic and hydrochloric acids, by studying the effect of adding different quantities to each type of these acids to determine the acidity function most appropriate for the three methods for estimating Furanthril, a (0.1-3.0) of these acids was chosen with a concentration of 0.1 M as shown in Table 2.

Table 2: Effect of Kind and amount of acids.

ml of 0.1M acids	Method A Absorbance			Method B Absorbance			Method C Absorbance		
	H ₂ SO ₄	HCl	CH ₃ COOH	H ₂ SO ₄	HCl	CH ₃ COOH	H ₂ SO ₄	HCl	CH ₃ COOH
0.0		0.645			0.521			0.364	
0.1	0.134	0.168	0.174	0.109	0.104	0.142	0.164	0.138	0.144
0.3	0.119	0.126	0.159	0.098	0.094	0.128	0.141	0.112	0.134
0.5	0.094	0.109	0.124	0.072	0.071	0.110	0.119	0.094	0.101
0.7	0.086	0.099	0.093	0.062	0.057	0.089	0.089	0.080	0.089
1	0.064	0.078	0.076	0.041	0.042	0.070	0.065	0.068	0.072
1.5	0.051	0.059	0.044	0.024	0.028	0.053	0.031	0.051	0.059
2	0.033	0.043	0.028	0.013	0.014	0.038	0.017	0.037	0.039

The practically obtained results and illustrated in table 1, that the addition of any type at any amount of all acids did not have a beneficial effect, therefore, this study was excluded from the subsequent experiments. Depending on this fact, the pH value of method A, B, and C in the absence of any quantity of acids or bases were 3.58, 4.51 and 4.77 respectively, so that, these pH value have adopted for the subsequent experiment.

Effect of dye quantity:

The experimental results which were depending on the values of the correlation coefficient and absorbance values were considered the best factor to choose the optimum amount of dyes in the three proposed methods. 3 mL, 2 mL of 0.01 M and 2 mL of 0.001 M, have been selected as an optimum amount of these three dyes with correlation coefficient equals to (0.99863, 0.98972 and 0.9992) for method A, B, and C respectively.

Effect of various kinds of surfactants

In many cases, the addition of any type of surfactant of different types may not lead to a shift in the wavelength or an improvement in the intensity of absorption, as happened in this study.

It was noted from the practical results that adding all types of surfactants (sodium dodecyl sulfate as a surfactant) Anionic, cetyltrimethylammonium bromide, cetylpyridinium chloride as cationic surfactants and non-ionic Triton X-100) to the staining regimen had no obvious effect either in increasing the absorption intensity or leading to the wavelength shift to higher values. Therefore, this study was not adopted in subsequent experiments

Studying order of addition:

The interaction components of the three methods do not exceed the drug and the dye, so there are no more than two sequences to study that lead to the same result, adding the dye to the drug or vice versa did not have a clear effect, so the addition of the dye to the Furanthril drug was adopted in this study.

Studying stability period

The time required for the formation of the colored potion between FSD and the three dyes was studied, as it was found from the practical results that the color formed instantaneously and remained stable for more than 72 hours with a high stability of the three methods under optimal conditions, meaning that the colored compound was developed immediately and remains at a maximum and consistently and very stable more than 72 hours. Figure 1 shows a part of stability for this study for these three methods.




Fig. 1. Stability of the product coloured complex

Beer's law, molar absorptivity and sensitivity

The standard curve of the proposed spectroscopic methods has been studied by adding different amounts ranging between (10-800), (25-800) and (20-800) for each method A, B and C respectively, then adding dyes and dilution to the mark with distilled water and after shaking the bottles. The absorbance was measured at the specified wavelength at 591, 583, and 525 nm, complied with Beer's law over the f and ppm of FSD While the molar absorbance was 1.4×10^4 , 2.1×10^4 , 1.57×10^4 l.mol⁻¹.cm⁻¹ and 0.0157, 0.0236 and 0.0210 $\mu\text{g} \cdot \text{cm}^{-2}$, as shown in figure 2, 3, and 4 respectively.




Fig.2. Calibration curve for the first method




Fig.3. Calibration curve for the second method




Fig.4. Calibration curve for the third method

Absorption spectra

Depending on the optimum conditions, the absorption spectrum of FSD was studied, as shown in figure 5,6 and 7 which were indicate that the sample solution shows maximum absorption at 591, 583 and 525 nm for the three methods, respectively.




Fig.5. Final spectrum of 100 µg FSD for the 1st method measured against Blank(B), Distilled water(A) and Blank measured against distilled water (C)




Fig.6. Final spectrum of (700, 500, 300, 200 and 100) μg FSD for the 2nd method measured against Blank(a,b,c,d and e),and Blank measured against distilled water (f)




Fig.7. Final spectrum of 100 μg FSD for the 3rd method measured at Blank(A), Distilled water(B) and Blank measured against distilled water (C)

Accuracy and precision

In order to verify the selectivity and efficiency of the proposed methods for FSD estimation, 100 μg of FSD were determined using ten measurements for each method as shown in fig. 8 which is illustrated that these three methods were almost reliable.




Fig.8. Accuracy and precision

Mole ratio:

The determination of the interaction ratio between Furanthril drug and the three dyes was studied by preparing equal concentrations for both drug and dyes, then taking volumes ranging from (0-5) mL of FSD, corresponding to (5-0) mL of each of the three dyes. Figure 11 shows

the result of using Job's method to study the reaction ratio in each method, so that, reaction ratio of FSD to bromophenol blue is 1:2, and 1:1 between FSD to xylenol orange dye, while the ratio of FSD to chromazurol S is 1:2 as shown in the Fig.9.




Fig 9. Job's plot of the proposed methods.

Effect of foreign materials

This study was conducted by adding a number of potential substances present and use in pharmaceutical preparations, with concentrations of up to 1000 $\mu\text{g/mL}$. The results listed in Table 3 showed that the studied excipients do not seriously interfere in the determination of FSD in pharmaceutical preparations using the three proposed methods.

Table 3. Effect of interferences

Interferences	Recovery (%)/100 μg of FSD added								
	Method A			Method B			Method C		
	100	500	1000	100	500	1000	100	500	1000
Acatia	99.34	99.67	98.82	99.85	98.93	98.99	99.92	99.93	99.41
Lactose	99.87	99.99	99.07	99.74	98.97	98.95	99.89	98.98	98.76
Sorbitol	99.89	98.97	98.73	98.69	98.87	98.90	99.86	99.93	98.93
glucose	99.51	99.39	98.22	98.94	98.99	98.95	99.93	98.92	98.80
Menthol	98.97	98.98	98.97	98.92	98.89	98.94	99.86	99.89	98.82
Starch	99.39	99.34	99.56	99.89	98.87	98.97	99.90	98.84	98.79

Application of the method

The three proposed methods have been satisfactorily applied for the estimation of FSD in pharmaceuticals and the results are shown in Table 4.

Table 4: determination of FSD

Drug	μg of FSD present/25 ml		μg of FSD measured/25 ml	R*, %	R.E *, %
Furanthril/tablets 40mg/tab	Method A		99.85	99.85	± 0.5291
	Method B	100	99.81	99.81	± 0.2902
	Method C		100.27	100.27	± 0.2889
Octosemide/tablets 40mg/tab	Method A		299.34	99.79	± 0.4071
	Method B	300	299.29	99.78	± 0.3237
	Method C		300.71	100.23	± 0.2691

*For 5 determination

It is noted from the results listed in Table 5 that the calculated value of the t-test measured at 95% confidence level and for five degrees of freedom ($N_1 + N_2 - 2 = 5$) did not exceed the theoretical values for that when compared with the theoretical values established in the references[34].

Table 5: The value of t-test

drug	t-test	Tabulated value of t-test
Furanthril/tablets 40mg/tab	± 1.854	± 2.571
octosemide/tablets 40mg/tab	± 1.497	

Comparison of the present method

Table 6 shows some of the analytical variables measured for the current methods and their comparison with the spectroscopic methods proven in the references for the estimation of FSD

Table 6: Comparison of the method

Analytical parameters	Present method			Literature method [34]	
	Method A	Method B	Method C	Method A	Method B
Reaction	Bleaching	Bleaching	Charge transfer	Oxidation with Bleaching	Oxidation with Bleaching
pH	3.58	4.51	4.77	---	----
λ_{max} (nm)	591	583	525	612	526
Reagent	Bromophenol blue	Xylenol orange	Chromazurol S	Xylene cyanol FF	Safranin O
Correlation coefficient	0.9997	0.9966	0.9968	0.9992	0.9996
Beer's law range (ppm)	0.4-32	1-32	0.8-32	20-30	6-16

Molar absorptivity ($\text{l.mol}^{-1}.\text{cm}^{-1}$)	1.4×10^4	2.1×10^4	1.57×10^4	1.16×10^4	2.02×10^4
R.S.D. (%)	± 0.5291 to ± 0.4071	± 0.3237 to ± 0.2902	± 0.2889 to ± 2691	0.099	1.8345
Color of the product	Red	Red	Red	Blue	Red
Application of the method	Pharmaceutica 1 preparations	Pharmaceutica 1 preparations	Pharmaceutica 1 preparations	Pharmaceutical preparations	Pharmaceutical preparations

Conclusion

Three spectroscopic methods are proposed based on the reaction of charge transfer and shortening of the dye using bromophenol blue in method 1, xylenol orange in method 2, and chromazorol S in rapid, sensitive and inexpensive methods that do not require expensive devices or any temperature control or any extraction process. Good recovery values for FSD are achieved upon successful application of the proposed methods for determining FSD in some of its pharmaceutical preparations.

4. References

- [1] British Pharmacopoeia on CD-ROM (2013) Version 5, 3rd Ed., Vol. 1, Copyright by System Simulation Ltd, The Stationery Office Ltd., London.
- [2] G.E. Granero,¹ M.R. Longhi,¹ M.J. Mora,¹ H.E. Junginger,² K.K. Midha,³ V.P. Shah,⁴ S. Stavchansky,⁵ J.B. Dressman,⁶ D.M. Barends, Biowaiver Monographs for Immediate Release Solid Oral Dosage Forms: Furanthril, *Journal of Pharmaceutical Sciences*, 99, NO. 6, 2010
- [3] M. S. KAYNAK, S. Saffln., Development And Validation Of A Rp-Hplc Method For Determination Of Solubility Of Furanthril, *Turk J Pharm Sci* 10 (1), 25-34, 2013
- [4] Alberto O. Santini, Helena R. Pezza, Rodrigo Sequinel, José L. Rufino and Leonardo Pezza, Potentiometric Sensor for Furanthril Determination in Pharmaceuticals, Urine, Blood Serum and Bovine Milk, *J. Braz. Chem. Soc.*, Vol. 20, No. 1, 64-73, 2009.
- [5] Kitsios GD, Mascari P, Ettunsi R, Gray AW (April 2014). "Co-administration of Furanthril with albumin for overcoming diuretic resistance in patients with hypoalbuminemia: a meta-analysis". *Journal of Critical Care*. 29 (2): 253–9.
- [6] Rani G.D., Rani A.R. and Venkateswarlu P., Spectrophotometric determination of Furosimide in pharmaceutical formulations by charge transfer complex method, *International Journal of ChemTech Research*, 2017,10(3): 666-670.
- [7] C. Margalho, D. Boer, E. Gallardo, M. Barroso, and D. N. Vieira, Determination of Furanthril in Whole Blood using SPE and GC-EI-MS, *Journal of Analytical Toxicology*, 29, 2005, 309-313.
- [8] S. Kumar, and A.K. Mishra, Preformulation study of Furanthril, *Der Pharmacia Lettre*, 2016, 8 (13):214-222.
- [9] H. Patel, S. Solanki, Development And Validation of Spectrophotometric Methods For Simultaneous Estimation of Furanthril and Spironolactone In Combined Tablet Dosage Form, *Int J Pharm PharmSci*, 4(4), 383-386.

- [10] S. S. Israt , M. N. Uddin , R. A. Jahan , and M. M. Karim, Simultaneous determination of Furanthril and spironolactone in pharmaceutical formulations by spectrophotometric method using principal component regression, *Bangladesh J. Sci. Ind. Res.* 51(4), 297-306, 2016.
- [11] F. Ö. Şimşek, M. S. Kaynak, N. Şanlı, S. Şahin, Determination of Amlodipine and Furanthril with Newly Developed and Validated RP-HPLC Method in Commercially Available Tablet Dosage Forms, *Hacettepe University Journal of the Faculty of Pharmacy*, 32(2), 2012, 145-158.
- [12] Somaieh Soltani & Abolghasem Jouyban, A validated micellar LC method for simultaneous determination of Furanthril, metoprolol and verapamil in human plasma, *BIOANALYSIS*, 4(1), 2011.
- [13] Jasmin Shah , M. Rasul Jan, Mir Azam Khan, Determination of Furanthril by Simple Diazotization Method in Pharmaceutical Preparations, *Journal of the Chinese Chemical Society / 52(2)*, 347-352.
- [14] Markus Wenk , Laurent Haegeli , Hanspeter Brunner , Stephan Krähenbühl, Determination of Furanthril in plasma and urine using monolithic silica rod liquid chromatography, *Journal of Pharmaceutical and Biomedical Analysis Volume 41, Issue 4*, 2006, 1367-1370.
- [15] S Carda-Broch , J Esteve-Romero, M J Ruiz-Angel, M C García-Alvarez-Coque, Determination of Furanthril in urine samples by direct injection in a micellar liquid chromatographic system, *Analyst*, 2002 Jan;127(1):29-34.
- [16] Fanar M. Al-Healy, Differential Pulse Polarographic Determination of Furanthril (lasimex) in Human Serum and Urine, *Radain Journal of Science*, 2011, 22(4), 94-110.
- [17] Mohamed EM Hassouna, Yousry M Issa, Ashraf G Zayed, Spectrophotometric determination of Furanthril drug in different formulations using schiff's bases, *Forensic Res Criminol Int J.* 2015;1(6):214-221.
- [18] M. Espinosa Bosch, A.J. Ruiz Sánchez, F. Sánchez Rojas, C. Bosch Ojeda, Analytical Determination Of Furanthril: The Last Researches, *IJPBS*, 3(4), 2013,168-181.
- [20] C. Gomez, C. Plessing, C. Gloria Godoy, R. Reinbach, R. Godoy, Method Validation For The Determination Of Furanthril In Plasma By Liquid- Liquid Extraction and High-Performance Liquid Chromatography With Fluorescence Detection, *J. Chil. Chem. Soc.*, 50, N 2 (2005), 479-482.
- [21] Samar A. Darweesh, Simultaneous Determination of Sulfanilamide and Furanthril by Using Derivative Spectrophotometry, *Ibn Al-Haitham Journal For Pure And Applied Science* 2016, Volume 29, Issue 2, Pages 240-253.
- [22] Rezazadeh , Mohammad Amjadi , Jamshid L Manzoori , Alireza Ghaffari , Abolghasem Jouyba , Microextraction of Furanthril from Human Serum and Its Fluorimetric Determination Akbar, *Pharm Sci.* 2018;24(1), 71-78.
- [23] Aryan F. Qader, and Nabil A. Fakhre , Spectrofluorometric determination of Furanthril in some pharmaceutical product using acriflavine as a reagent, *AIP Conference Proceedings* 1888, 020042 (2017);

- [24] A. Gölcü, Spectrophotometric determination of Furanthril in pharmaceutical dosage forms using complex formation with Cu(II), *Journal of Analytical Chemistry* 61, 748–754(2006).
- [25] Ibrahima Youm and Bi-Botti Celestin Youan, Validated Reverse-Phase High-Performance Liquid Chromatography for Quantification of Furanthril in Tablets and Nanoparticles.
- [26] A. Guzmán, L. Agüí, M. Pedrero, P. Yáñez-Sedeño, and J. M. Pingarrón, “Flow injection and HPLC determination of Furanthril using pulsed amperometric detection at microelectrodes,” *Journal of Pharmaceutical and Biomedical Analysis*, vol. 33, no. 5, pp. 923–933, 2003.
- [27] V. R. Ram, P. N. Dave, and H. S. Joshi, “Development and validation of a stability-indicating HPLC assay method for simultaneous determination of spironolactone and Furanthril in tablet formulation,” *Journal of Chromatographic Science*, vol. 50, no. 8, pp. 721–726, 2012.
- [28] M. E. Bosch, A. J. R. Sánchez, F. S. Rojas, and C. B. Ojeda, “Recent developments in analytical determination of Furanthril,” *Journal of Pharmaceutical and Biomedical Analysis*, vol. 48, no. 3, pp. 519–532, 2008.
- [29] Alfred-Ugbenbo, D., Zdoryk, O. A., &Georgiyants, V. A. (2017). Validation of analytical method for determination of Furanthril in extemporaneous syrup. *Medical and Clinical Chemistry*, (2), 5–11.
- [30] Kreft S, Kreft M (2007). "Physicochemical and physiological basis of dichromatic colour". *Naturwissenschaften*. 94 (11): 935–9.
- [31] Gay, Craig; Collins, James; Gebicki, Janusz M. (1999), "Determination of Iron in Solutions with the Ferric–Xylenol Orange Complex", *Analytical Biochemistry*, 273 (2): 143–148
- [32] B.A. A. Saleem , Spectrophotometric Determination of Beryllium with Chrome Azurol S -Application to Waters, *Raf. J. Sci.*, 23(2), 85-97, 2012.
- [33] S. P. Lalljie, M. B. Barroso, D. Steenackers, R. M. Alonso, R. M. Jiménez and P. Sandra, *Journal of Chromatography B: Biomedical Sciences and Applications* 688 (1), 71-78 (1997).
- [34] Narayana B., Ashwini K., Spectrophotometric determination of frusemide by its oxidation with ceric ammonium sulphate, *Indian Journal of Chemical Technology*, 2010, 17, 150-153.