

Synthesis, Characterization and Antimicrobial Activity of Mixed Ligand Complexes of N-(2-hydroxy-1-naphthylidene)-4-nitroaniline and N-(2-hydroxy-3-methoxybenzylidene)-4-methylaniline with Co(II), Ni(II), Cu(II) and Mn(II) ions.

Dr. Madhukar A. Badgujar

Department of Chemistry, Sheth J.N. Paliwala College Pali- Sudhagad, Raigad(M.S)

email- bmbadgujar@yahoo.com

Abstract:

The mixed ligand complexes of Co(II), Ni(II), Cu(II) and Mn(II) with Schiff bases N-(2-hydroxy-1-naphthylidene)-4-nitroaniline (L^1H) and N-(2-hydroxy-3-methoxy benzylidene)-4-methylaniline (L^2H) have been synthesized and characterized. The resulting complexes were characterized by elemental analysis, thermogravimetric analysis, magnetic moment measurements, conductivity measurements, 1H NMR, IR, UV-visible and ESR spectral studies. The Schiff bases acts as bidentate monobasic ligands, coordinating through deprotonated phenolic oxygen and azomethine nitrogen atoms. The complexes are non-electrolytic in DMSO. The presence of the two coordinated water molecules in these complexes was indicated by IR spectra and thermogravimetric analysis of the complexes. From the analytical and spectral data the stoichiometry of these complexes have been found to be $[M(L^1)(L^2)(H_2O)_2]$ {where M = Co(II), Ni(II), Cu(II) and Mn(II)}. It is found that Co(II), Ni(II), Cu(II) and Mn(II) complexes exhibited octahedral geometry. The antimicrobial activities of ligands and their mixed ligand complexes were screened by Disc Diffusion method. It was found that the metal complexes have higher antimicrobial activity than the free ligand.

Keywords: Schiff bases, Mixed ligand, Metal complexes, Antimicrobial activity.

Introduction

Mixed ligand complexes play an important role in biological processes like activation of enzymes by metals. Such complexes are useful in the storage and transport of active substances through membranes. Schiff bases play important roles in coordination chemistry as they easily form stable complexes with most transition metal ions [1,2]. The present course of study was initiated to understand the characteristic nature and applications of the biologically active metal complexes of nitrogen and oxygen donor ligands. By considering all these applications, in this paper we are reporting the synthesis and characterization of series of Mn(II), Co(II), Ni(II) and Cu(II) complexes obtained from N-(2-hydroxy-1-naphthylidene)-4-nitroaniline and N-(2-hydroxy-3-methoxybenzylidene)-4-methylaniline as ligands. Many biologically important Schiff bases have been reported in the literature possessing, antimicrobial, antibacterial, antifungal, anti-inflammatory, anticonvulsant, antitumor and anti HIV activities [3-8].

Experimental

All the chemicals used in the present work were of analytical grade. The metal salts used were in their hydrated form.

IJFANS INTERNATIONAL JOURNAL OF FOOD AND NUTRITIONAL SCIENCES

ISSN PRINT 2319 1775 Online 2320 7876

Research Paper © 2012 IJFANS. All Rights Reserved. UGC CARE Listed (Group -I) Journal Volume 11, Iss 08, 2022

Synthesis of ligands

The ligands (L^1H and L^2H) were prepared by the condensation of the corresponding aldehyde and amine in 1:1 molar ratio by refluxing in distilled methanol [9].

Preparation of complexes

A metal(II) chloride (1.0 mmol.) dissolved in water (2 mL) was added slowly with constant stirring to a solution of N-(2-hydroxy-1-naphthylidene)-4-nitroaniline (L^1H) (1.0 mmol.) and N-(2-hydroxybenzylidene)-4-methylaniline (L^2H) (1.0 mmol.) in ethanol (100 mL). The pH of the solution was slowly raised to obtain the appropriate pH for the formation of the complex by the drop wise addition of 0.1 N sodium hydroxide solution. The resulting mixture was stirred for 2-3 hour. The product was digested on a hot water bath for half an hour. The complex thus formed was filtered out, washed with aqueous alcohol followed by distilled water and dried in vacuum over fused $CaCl_2$. Similarly all the complexes were prepared. The metal estimations were carried out by standard Methods [10]. Elemental analysis was performed at SAIF, IIT Bombay, Mumbai. The conductance was measured in DMSO solvent on an Equiptronic EQ-660 conductivity meter. The magnetic susceptibility measurements at room temperature were made on Gouy balance using $HgCo(NCS)_4$ as calibrant. The 1H NMR spectrum of the ligands and Zn(II) complex was recorded on Varian Mercury plus 400 MHz spectrometer using TMS as internal standard and (D_6) DMSO as a solvent. The IR spectra of ligands and their complexes were recorded on a Shimadzu FTIR – 8400S spectrometer in KBr pellets in the range of 4000-350 cm^{-1} . UV-Visible spectra were recorded on a Jasco V-530 UV-Visible spectrometer in the range 200-1100 nm. Electron spin resonance spectra of powdered Cu(II) complex was recorded on Varian E-112x-band ESR spectrometer using TCNE as 'g' marker ($g = 2.0027$) at room temperature. Thermal analysis of the metal complexes was carried out using Mettler Toledo Star^e SW 7.01 thermal analyzer in an inert atmosphere of nitrogen. The antimicrobial activities of ligands and their mixed ligand complexes were screened by Disc Diffusion method.

Results and Discussion

The elemental analysis shown in the Table 1 indicates that, all the metal complexes have 1:1:1 stoichiometry with respect to L^1H : L^2H : M {where M = Co(II), Ni(II), Cu(II) and Mn(II)} and are dark colored amorphous substances, soluble in DMF and DMSO. The molar conductance values (0.012 - 0.015 $ohm^{-1} mol^{-1} cm^2$) of 10^{-3} M solution in DMSO indicate that the metal complexes are non-electrolytic in nature.

Table 1. Analytical and Physical data of Schiff bases L^1H , L^2H and their mixed ligand metal complexes.

Ligand / complex, Molecular Formula	Mol. Wt.	M.P / Decomp. Temp. °C	Analysis % Found (Calculated)				Molar Conductance (λm) $Ohm^{-1} cm^2 mol^{-1}$
			C	H	N	M	
L^1H $C_{17}H_{12}N_2O_3$	292.29	138	69.65 (69.86)	4.34 (4.14)	9.71 (9.58)	-	-

IJFANS INTERNATIONAL JOURNAL OF FOOD AND NUTRITIONAL SCIENCES

ISSN PRINT 2319 1775 Online 2320 7876

Research Paper © 2012 IJFANS. All Rights Reserved. UGC CARE Listed (Group -I) Journal Volume 11, Iss 08, 2022

L^2H $C_{14}H_{12}NO$	225.29	91	79.75 (79.97)	6.63 (6.71)	6.12 (6.21)	-	-
$[Co(L^1)(L^2)(H_2O)_2]$ $C_{32}H_{29}N_3O_6Co$	610.25	>250	62.67 (62.95)	4.71 (4.79)	6.67 (6.88)	9.78 (9.65)	0.018
$[Ni(L^1)(L^2)(H_2O)_2]$ $C_{32}H_{29}N_3O_6Ni$	610.28	>250	62.81 (62.98)	4.91 (4.79)	6.78 (6.890)	9.75 (9.62)	0.012
$[Cu(L^1)(L^2)(H_2O)_2]$ $C_{32}H_{29}N_3O_6Cu$	615.14	>250	62.57 (62.48)	4.87 (4.75)	6.67 (6.83)	10.42 (10.33)	0.014
$[Mn(L^1)(L^2)(H_2O)_2]$ $C_{32}H_{29}N_3O_6Zn$	617.00	>250	62.11 (62.29)	4.92 (4.74)	6.67 (6.81)	10.51 (10.60)	0.015

Electronic spectra and Magnetic moment

The electronic spectral studies of Mixed Ligand Complexes of Co(II), Ni(II), Cu(II) and Zn(II) were carried out in DMSO solution. The absorption spectrum of the Co(II) complex shows bands at $\sim 9523.81 \text{ cm}^{-1}$ ($\epsilon \sim 27 \text{ L mol}^{-1} \text{ cm}^{-1}$) and $\sim 19047.62 \text{ cm}^{-1}$ ($\epsilon \sim 60 \text{ L mol}^{-1} \text{ cm}^{-1}$) attributed to ${}^4T_{1g}(F) \rightarrow {}^4T_{2g}(F)$ (ν_1) and ${}^4T_{1g}(F) \rightarrow {}^4T_{1g}(P)$ (ν_3) transitions respectively in an octahedral field [11]. The spectral parameters of Co(II) complex are as follows [12]: $Dq = 1064.85 \text{ cm}^{-1}$, $B = 709.9 \text{ cm}^{-1}$, $\beta = 0.731$ and $\beta^0 = 26.9 \%$. The reduction of Racah parameter from the free ion value 971 cm^{-1} and β^0 value of 26.9% testify the presence of considerable covalence in the complex [13]. The Co(II) complex has magnetic moment 5.14 B.M. also suggest an octahedral geometry. Ni(II) complex exhibits two electronic spectral bands at $\sim 10504.20 \text{ cm}^{-1}$ ($\epsilon \sim 30 \text{ L mol}^{-1} \text{ cm}^{-1}$) and $\sim 16393.44 \text{ cm}^{-1}$ ($\epsilon \sim 87 \text{ L mol}^{-1} \text{ cm}^{-1}$) which can be assigned to ${}^3A_{2g}(F) \rightarrow {}^3T_{2g}(F)$ (ν_1) and ${}^3A_{2g}(F) \rightarrow {}^3T_{1g}(F)$ (ν_2) transitions in an octahedral field [14]. The transition due to ${}^3A_{2g}(F) \rightarrow {}^3T_{1g}(P)$ (ν_3) was completely obscured by the intense intra ligand transition band. The $\nu_2 : \nu_1$ ratio is 1.56 , which is in the usual range reported for an octahedral Ni(II) complexes [15]. The spectral parameters of the Ni(II) complex are as follows [12]: $Dq = 1050.23 \text{ cm}^{-1}$, $B = 724.30 \text{ cm}^{-1}$, $\beta = 0.703$ and $\beta^0 = 29.7 \%$. The reduction of Racah parameter from the free ion value 1030 cm^{-1} and β^0 value of 29.7% confirms the presence of considerable covalence in the complex [13]. The Ni(II) complex has magnetic moment 3.47 B.M. also suggest an octahedral geometry. The Cu(II) complex exhibit broad band centered at $\sim 14285.71 \text{ cm}^{-1}$ ($\epsilon \sim 120 \text{ L mol}^{-1} \text{ cm}^{-1}$) mainly due to ${}^2E_g \rightarrow {}^2T_{2g}$ transition suggesting the distorted octahedral geometry [16]. The observed magnetic moment value for Cu(II) complex is 1.81 B.M. suggestive of distorted octahedral nature for the complex [16]. Zn(II) complex does not exhibit any characteristic d-d transitions and also found to be diamagnetic in nature.

Infrared spectra

The important infrared frequencies exhibited by the ligands L^1H and L^2H and their mixed ligand metal complexes are given in the Table 2. Infrared spectra of the schiff bases L^1H and L^2H show a broad band centered at around 3437 cm^{-1} and 3447 cm^{-1} due to $\nu(\text{OH})$ of the phenolic hydroxyl group respectively in free ligands, which disappeared in spectra of their complexes indicating probably the coordination through phenolic oxygen moiety. The schiff bases L^1H and L^2H show a

IJFANS INTERNATIONAL JOURNAL OF FOOD AND NUTRITIONAL SCIENCES

ISSN PRINT 2319 1775 Online 2320 7876

Research Paper © 2012 IJFANS. All Rights Reserved. UGC CARE Listed (Group -I) Journal Volume 11, Iss 08, 2022

medium intensity band at around 1335 cm^{-1} and 1279 cm^{-1} due to $\nu(\text{C-O})$ (phenolic) is shifted to higher region at around $1381\text{-}1399\text{ cm}^{-1}$ and $1359\text{-}1369\text{ cm}^{-1}$ respectively indicating the coordination through the phenolic oxygen atoms [14,15,17]. The IR spectra of the schiff bases L^1H and L^2H exhibit a strong band at 1622 cm^{-1} and 1613 cm^{-1} due to $\nu(\text{C=N})$ (azomethine) which has been shifted towards lower region at around $1613\text{-}1616\text{ cm}^{-1}$ and $1601\text{-}1602\text{ cm}^{-1}$ respectively in the spectra of complexes indicating the participation of the azomethine groups in the complex formation [17,18]. The spectra of the complexes show a broad diffused bands in the region at around $3150\text{-}3700\text{ cm}^{-1}$, strong bands at $1534\text{-}1537\text{ cm}^{-1}$ and week intensity bands at $829\text{-}836\text{ cm}^{-1}$ due to $\nu(\text{OH})$, $\delta(\text{OH})$ and $\rho_r(\text{OH})$ respectively of the coordinated water molecules [14,15]. The coordination through nitrogen of azomethine and oxygen of (C-O) group of ligands are further evidenced by the appearance of non-ligand bands in the complexes at around $503\text{-}557\text{ cm}^{-1}$ and $416\text{-}463\text{ cm}^{-1}$ are due to $\nu(\text{M-O})$ and $\nu(\text{M-N})$ vibrations respectively [17,18].

Table 2. Characteristic IR bands of the Schiff Bases L^1H , L^2H and their mixed ligand metal complexes.

Schiff base / Complex	IR bands (cm^{-1})							
	ν_{OH} (phenolic)	ν_{OH} (H_2O)	$\nu_{\text{C=N}}$	δ_{OH} (H_2O)	$\nu_{\text{C-O}}$ (phenolic)	ρ_{rOH} (H_2O)	$\nu_{\text{M-O}}$	$\nu_{\text{M-N}}$
L^1H	3437	-	1622	-	1335	-	-	-
L^2H	3447	-	1613	-	1279	-	-	-
$[\text{Co}(\text{L}^1)(\text{L}^2)(\text{H}_2\text{O})_2]$	-	3150-3650	1614, 1601	1535	1381, 1359	834	550, 511	457, 416
$[\text{Ni}(\text{L}^1)(\text{L}^2)(\text{H}_2\text{O})_2]$	-	3150-3700	1615, 1602	1534	1399, 1369	831	554, 503	459, 418
$[\text{Cu}(\text{L}^1)(\text{L}^2)(\text{H}_2\text{O})_2]$	-	3200-3650	1613, 1601	1536	1383, 1360	829	555, 508	461, 421
$[\text{Zn}(\text{L}^1)(\text{L}^2)(\text{H}_2\text{O})_2]$	-	3150-3700	1616, 1602	1537	1381, 1362	836	557, 510	463, 422

^1H NMR spectra

The ^1H NMR spectrum of the ligand (L^1H) showed singlet at $15.217\text{ }\delta$ (s, 1H, OH) due to proton of phenolic -OH group. The azomethine proton have resonated as singlet at $9.682\text{ }\delta$ (s, 1H, CH=N). The ten aromatic protons have resonated in the region $6.594\text{-}8.545\text{ }\delta$ (m, 10H, Ar-H) as multiplet. The ^1H NMR spectrum of the ligand (L^2H) showed singlet at $13.291\text{ }\delta$ (s, 1H, OH) due to proton of phenolic -OH group. The azomethine proton have resonated as singlet at $8.77\text{ }\delta$ (s, 1H, CH=N). The seven aromatic protons have resonated in the region $6.956\text{ - }7.622\text{ }\delta$ (m, 7H,

IJFANS INTERNATIONAL JOURNAL OF FOOD AND NUTRITIONAL SCIENCES

ISSN PRINT 2319 1775 Online 2320 7876

Research Paper © 2012 IJFANS. All Rights Reserved. UGC CARE Listed (Group -I) Journal Volume 11, Iss 08, 2022

Ar-H) as multiplet. The six protons of two methyl groups resonated as two singlets at 2.198 δ (s, 3H, Ar-CH₃) and 2.251 δ (s, 3H, Ar-CH₃).

The ¹H NMR spectrum of the [Zn(L¹)(L²)(H₂O)₂] complex showed two singlet at 9.281 δ (s, 1H, CH=N) and 8.511 δ (s, 1H, CH=N) due to two protons of the two azomethine groups. The seventeen aromatic protons have resonated in the region 6.303-8.306 δ (m, 17H, Ar-H) as multiplet. The six protons of two methyl groups resonated as two singlets at 1.975 δ (s, 3H, Ar-CH₃) and 2.115 δ (s, 3H, Ar-CH₃). The proton of phenolic –OH group which was observed at 15.217 δ (s, 1H, OH) and 13.291 δ (s, 1H, OH) for Schiff bases L¹H and L²H respectively has disappeared in Zn(II) complex, indicating coordination through phenolic oxygen moiety to the metal ion via deprotonation [15,17].

Thermogravimetric analysis

The dynamic TGA with the percentage mass loss at different steps have been recorded. The elimination of lattice and coordinated water molecules take place in the first step. The Co(II), Ni(II), Cu(II) and Zn(II) complexes lose their weight in the temperature range ~ 125-290 °C, 130-285 °C, 125-285 °C and 130-290 °C respectively corresponding to two coordinated water molecules with an endothermic peak in DTA curve. After the total loss of water, the organic moiety decomposes on further increment of temperature. The complete decomposition of ligands occurs at ~ 450-650 °C and the observed residue corresponds to the respective oxide. The occurrence of endothermic peaks in DTA curves and elimination of the two water molecules at comparatively higher temperature unambiguously confirm our earlier observation based upon the IR spectrum that the water molecules are coordinated in the metal complexes [11,14].

ESR spectra

The ESR spectrum of the powdered sample of the Cu(II) complex was recorded at room temperature. The observed value of ESR spectral parameters for the [Cu(L¹)(L²)(H₂O)₂] complex are $g_{\parallel} = 2.137$, $g_{\perp} = 2.083$ and $G = 1.667$. The observed g_{\parallel} value is less than 2.3 in agreement with the covalent character of the metal – ligand bond [13]. The trend $g_{\parallel} > g_{\perp} > g_e$ (2.0023) observed for this complex shows that the unpaired electron is localized in $d_{x^2-y^2}$ orbital of Cu(II) ion and the spectral features are characteristic of axial symmetry; tetragonal elongated structure may be assumed for this Cu(II) complex [13,19]. The anisotropic G values have been calculated by using the equation $G = (g_{\parallel} - 2.002) / (g_{\perp} - 2.002)$. If the G value is less than 4.0, the ligand forming the copper(II) complex is regarded as strong field ligand while a value of G greater than 4.0 indicates weak field ligand. As $G = 1.667$ for the present complex indicates that the ligands L¹H and L²H are strong field ligands and the metal-ligand bonding in this complex is covalent [13].

Antimicrobial activity

Synthesized Schiff bases and their corresponding mixed ligand metal complexes were screened against microorganisms such as *E.coli*, *S.aurios*, *B.subtilis*, and *S.typhi* to assess their potential as antimicrobial agent by Disc Diffusion method. The zones of inhibition based upon zone size

IJFANS INTERNATIONAL JOURNAL OF FOOD AND NUTRITIONAL SCIENCES


ISSN PRINT 2319 1775 Online 2320 7876

Research Paper © 2012 IJFANS. All Rights Reserved. UGC CARE Listed (Group -I) Journal Volume 11, Iss 08, 2022

around the discs were measured. The measured zone of inhibition against the growth of various microorganisms is listed in Table 3. It is found that the metal complexes have higher antimicrobial activity than the free ligand. Hence complexation increases the antimicrobial activity [15,17].

Table 3. Antimicrobial activity data of the ligands L^1H , L^2H and their mixed ligand metal complexes.

Schiff base / Complex	Zone of inhibition (mm)			
	S.aureus	B.subtilis	E.coli	S.typhi
L^1H	2.7	2.9	0.0	2.5
L^2H	0.0	2.5	2.7	2.5
$[Co(L^1)(L^2)(H_2O)_2]$	8.0	7.5	6.5	4.1
$[Ni(L^1)(L^2)(H_2O)_2]$	6.3	7.7	6.5	3.9
$[Cu(L^1)(L^2)(H_2O)_2]$	3.6	6.5	7.1	4.8
$[Zn(L^1)(L^2)(H_2O)_2]$	8.0	9.3	4.5	6.3



M = Co(II), Ni(II), Cu(II) and Zn(II).

Figure 1. Proposed structure for the complexes

Conclusion

The elemental analysis, magnetic susceptibility, electronic, 1H NMR, IR and ESR spectral observations suggest the octahedral geometry for the Co(II), Ni(II), Cu(II) and Mn(II) complexes and exhibit coordination number six. The general structure of the complexes is shown in figure 1. The Schiff's bases do show some antimicrobial activity to certain extent but their complexes exhibit comparatively greater amount of activity against the microorganisms.

References

- [1] Spinu C., Pleniceanu M. and Tigae C., Biologically Active Transition Metal Chelates with a 2-Thiophenecarboxaldehyde-Derived Schiff Base: Synthesis, Characterization, and Antibacterial Properties, *Turk. J. Chem.*, 2008, 32, 487.
- [2] Clarke B., Clarke N., Cunningham D., Higgins T., McArdle P., Cholchu M.N. and O'Gara M., Transition-metal Schiff-base complexes as ligands in tin chemistry. Part 7. Reactions of organotin(IV) Lewis acids with $[M(L)]_2$ [M-Ni, Cu and Zn; H₂L-N,N-bis(3-methoxysalicylidene)benzene-1,3-diamine and its -1,4-diamine analog], *J. Organometallic Chem.*, 1998, 55, 559.
- [3] Pandeya S.N., Sriram D., Nath G. and Clercq E. De, Synthesis, antibacterial, antifungal and antiviral activity evaluation of some new bis-Schiff bases of isatin and their derivatives, *Pharm. Acta Helv.*, 1999, 74, 11.
- [4] Pandeya S.N., Sriram D., Nath G. and Clercq E. de, Synthesis, antibacterial, antifungal and anti-HIV evaluation of Schiff and Mannich bases of isatin and its derivatives with triazole, *Arzneimittel Forsch*, 2000, 50, 55.
- [5] Singh W.M. and Dash B.C., Synthesis of some new schiff bases containing thiazole and oxazole nuclei and their fungicidal activity, *Pesticides*, 1988, 22, 33.
- [6] Kelley J.L., Linn J.A., Bankston D.D., Burchall C.J., Soroko F.E. and Cooper B.R., 8-Amino-3-benzyl-1,2,4-triazolo[4,3- α]pyrazines. Synthesis and anticonvulsant activity, *J. Med. Chem.*, 1995, 38, 3676.
- [7] G. Turan-Zitouni, Z.A. Kaplancikli, A. Ozdemir, P. Chevallet, Studies on 1,2,4-triazole derivatives as potential anti-inflammatory agents, *Arch. Pharm. Chem. Life Sci.*, 2007, 340, 586.
- [8] Tarafder M.T.H., Kasbollah A., Saravanan N., Crouse K.A., Ali A.M. and Oo K.T., S-methyldithiocarbamate and its schiff bases: Evaluation of bondings and biological properties, *J. Biochem. Mol. BioI. Biophys.*, 2002, 6, 85.
- [9] Abd-Elzar M.M., Spectroscopic characterization of some tetradentate Schiff bases and their complexes with nickel, copper and zinc, *J. Chin. Chem. Soc.*, 2001, 48, 153.
- [10] Vogel A.I., Text Book of Quantitative Practical Inorganic Chemistry, ELBS, 1984.
- [11] Dash D.C., Meher F.M., Monhanty P.C. and Nanda J., Complexes of 3-Benzylideneimino-2-thio-hydantoin and 3-p-Anisylidenemino-2-thiohydantoin with Mn(II), Co(II), Ni(II) and Cu(II), *Indian J. Chem.*, 1987, 26(A), 698-701.
- [12] Liver A.B.P., Inorganic electronic spectroscopy. Elsevier, New York, 1984.
- [13] Dutta R. L. and Syamal A., Elements of Magnatochemistry, 2nd Ed., East west press, New Delhi, 1996.
- [14] Makode J.T. and Aswar A.S., Synthesis Characterization, biological and thermal properties of some new Schiff base complexes derived from 2-hydroxy-5-chloroacetophenone and S-methyldithiocarbamate, *Indian J. Chem.*, 2004, 43(A), 2120-2125.

IJFANS INTERNATIONAL JOURNAL OF FOOD AND NUTRITIONAL SCIENCES

ISSN PRINT 2319 1775 Online 2320 7876

Research Paper © 2012 IJFANS. All Rights Reserved. UGC CARE Listed (Group -I) Journal Volume 11, Iss 08, 2022

- [15] Ravanasiddappa M., Sureshg T., Syed K., Radhavendray S. C. Basavaraja C. and Angadi S.D., Transition Metal Complexes of 1, 4(2'-Hydroxyphenyl-1-yl) di-iminoazine: Synthesis, Characterization and Antimicrobial Studies, E-J. Chem., 2008, 5(2), 395-403.
- [16] Shriodkar S.G., Mane P.S. and Chondhekar T.K., Synthesis and fungitoxic studies of Mn(II), Co(II), Ni(II) and Cu(II) with some heterocyclic Schiff base ligands, Indian J. Chem., 2001, 40A, 1114-1117.
- [17] Reddy V., Patil N. and Angadi S.D., Synthesis, Characterization and Antimicrobial Activity of Cu(II), Co(II) and Ni(II) Complexes with O, N, and S Donor Ligands, E-J. Chem., 2008, 5(3), 577-583.
- [18] Nakamoto K., Infrared Spectra of inorganic and coordination Compounds, John Wiley, New York, 1970.
- [19] Thakkar N.V and Bootwala S.Z., Synthesis and characterization of binuclear metal complexes derived from some isonitrosoacetophenones and benzidine, Indian J. Chem. 1995, 34(A), 370-374.