

Phytochemical Screening And Separation Of Various Extracts Of *Cyanobacterium Nostoc Muscorum*

Sarita Dubey^{1*}

¹Department of Biotechnology and Bioinformatics Centre, Barkatullah University, Bhopal

Dr Sarita Khare²

²Department of Zoology, Government MLB Girls PG College, Bhopal

Dr Ankita Soni³

³Department of Zoology, SMS Government Model Science PG College, Gwalior

Dr Santosh Bhargava⁴

⁴Division of Microbiology, Department of Botany, Government Motilal Science College, Bhopal

***Corresponding Author: Dr Ankita Soni**

*Department of Biotechnology and Bioinformatics Centre, Barkatullah University, Bhopal

Email address: adr445777@gmail.com

Abstract:

Cyanobacteria is a free-living microorganism surviving in both terrestrial and fresh water habitat. Various strain of cyanobacteria is well known for the production of intracellular and extracellular metabolites like antimycobacterial, antifungal, antiviral and antialgal activity. In the present study, we performed phytochemical screening of *Cyanobacteria Nostoc muscorum* to extract and found the presence of alkaloids, flavonoids, saponins, and glycosides; and thin layer chromatography (TLC) of cyanobacteria *Nostoc muscorum*. Afterward, Thin layer chromatography of extract obtained by using different solvents of non-diazotrophically and diazotrophically grown culture of the *Cyanobacterium Nostoc muscorum* were performed.

Keywords: Cyanobacteria, *Nostoc muscorum*, Phytochemical screening.

1. INTRODUCTION

Cyanobacteria are one of the important organisms having some most important functions like nitrogen fixation, gram-negative and photoautotrophic nature. Cyanobacteria are known for their diversity in terms of morphological properties. Cyanobacteria are also known as blue-green algae and include a highly diverse group of photosynthetic prokaryotic microorganisms. They are widely distributed in nature and can be found in most terrestrial and freshwater habitats (Potts, 2002). In addition, Cyanobacteria can be found in every light Exposed habitat on earth and are a group of gram-negative eubacteria capable of oxygenic photosynthesis same as higher plants and thus they are considered to be the ancestors of higher chloroplast (Rodriguez et al., 2005). Recently, micro algae have become commercially important because of novel compounds and potential medicinal value. Microalgae are known to secrete vitamins, amino acids, siderophores, simple carbon hydrates, and other compounds that are essential or support the growth of other microorganisms. In recent years, microalgae have become an economic source of new drugs and commercially important compounds (Meeting and Pyne 1986). The biologically active compounds isolated from microalgae are known to show antibacterial, antiviral, antifungal, enzyme-inhibiting, immunostimulant, cytotoxic and antiplasmodial activities (Ghasemi et al. 2004). The secondary metabolites produced by Cyanobacteria are rich sources of

novel bioactive compounds applicable for the production of medicines and agriculturally important chemicals. In the natural environment, Cyanobacteria secrete some of the extracellular metabolites, function as toxins or allelochemicals (Pflugmacher 2002). The antifungal compound namely cryptophycin 1 is known to show potential activity against a large number of agriculturally important fungi (Biondi et al. 2004). The *Cyanobacterium Nostoc spongiaeforme* TISTR 8169 synthesizes and releases a violet pigment known as Nostocine A into the medium. Bioactive metabolites from cyanobacteria are also of biotechnological interest, particularly in the field of pharmaceutical industries. Cyanobacteria grow and multiply in minimal culture media on a mass scale. The growth condition can be manipulated to achieve optimum production of desirable bioactive compounds (Dahms et al. 2006). Many species of cyanobacteria are known to produce intracellular and extracellular metabolites. These metabolites show diverse biological activities and are known to inhibit microbial growth (Noaman et al. 2004). The *Cyanobacterium thormidium* has a broad range of antimicrobial activity (Fish and Codd 1994). So that the present study was performed for phytochemical screening of *Cyanobacteria Nostoc muscorum*. After the extraction of different secondary metabolites: alkaloids, flavonoids, saponins and glycosides were extracted and furthermore TLC were performed.

2. MATERIALS AND METHODS

In the present section of the work, the experiment organism was *Nostoc muscorum*. This species of cyanobacteria is a filamentous gram-negative, green brown color and heterocyst forming cyanobacteria. This Cyanobacteria were cultured in chu no.10 as described by Gerloff et al 1950. The ideal pH for the growth of this organism is 7.0-8.5. The culture was maintained at 28°C with the intensity of 14.40 w/m² provided by a white fluorescent tube with a light/dark cycle of 16/8 hours.

For the isolation of bioactive compounds, one-week old *Nostoc muscorum* (fully grown) was centrifuged and the pellet was dried in a hot air oven at 60°C. It was separated using a Soxhlet extractor, by using Chloroform, Acetone, Ethanol, Butanol, and Methanol. Chemical test for the screening and identification of bioactive chemicals like alkaloids, carbohydrates, glycosides, phenolic compounds, amino acids, carbohydrates, saponins, and protein were performed.

The absorption effect was studied on the basis of Thin layer chromatography. In this procedure, the form of mobile phase containing the dissolved solutes moves over the stationary phase. Various solvent extract was subjected to thin layer chromatography. In the matching through the chamber with different solvent systems (toluene: ethyl acetate: formic acid in 5:4:1 ratio) solvent system was used. After pre-saturation with mobile phase for 20 min, the development was achieved. So that, as the run plates were dried, they were sprayed with freshly prepared iodine reagents, to detect the bands on the TLC plates. The movement of the active compounds was expressed by its retention factors (R_f), and values were calculated for different samples.

$$R_f = \frac{\text{Distance traveled by solute}}{\text{Distance traveled by the solvent}}$$

3. RESULTS AND DISCUSSION

To Obtain the percentage of yield of extraction is a very important finding in phytochemical extraction. So, firstly the percentage yield of different extracts of the *Cyanobacterium Nostoc muscorum* were found (table no.1). We found 2.7%, 7.3%, 5.5%, 5.3%, and 5.9% yield in non-diazotrophic cultures (Medium containing 1nM NH₄Cl) using different solvents n-butanol, chloroform, acetone, ethanol, and methanol, respectively. Chloroform /Methanol extract of non-diazotrophic grown culture exhibited maximum percentage yield in comparison to all solvents. When compared with non-diazotrophic, diazotrophic extract were also showed similar trends of percentage

yield that is 1.2%, 5.5%, 4.2%, and 4.9% using n-butanol, chloroform, acetone, ethanol, and methanol extract, respectively. Chloroform/Methanol extract of diazotrophic–grown culture also exhibited maximum percentage yield in comparison to all solvents. This varying amount of percentage yield is because of the different polarity of solvent used for extraction.

Extract	Percentage yield (%)				
	n-Butanol	Chloroform	Acetone	Ethanol	Methanol
Diazotrophic grown culture	1.2 ±0.11	5.5 ± 0.48	4.2 ±0.40	4.6 ±0.39	4.9 ±0.44
Medium containing 5mM KNO ₂	1.8 ±0.15	6.6 ±0.52	4.5 ±0.41	4.9 ±0.41	5.2 ±0.48
Medium containing 5mM KNO ₃	2.2 ±0.19	6.9 ±0.55	5.1 ±0.49	5.1 ±0.48	5.4 ±0.52
Medium containing 1mM NH ₄ Cl	2.7 ±0.20	7.3 ±0.66	5.5 ±0.51	5.3 ±0.51	5.9 ±0.55

Table 1: The percentage yield of different extracts of Cyanobacterium *Nostoc muscorum* (Non–diazotrophs, and diazotrophs).

Phytochemical analysis

When a small amount of dried extract was subjected to phytochemical analysis to test for the presence of glycoside, saponin, tannins, flavonoids and steroids, the results are presented in table no.2. The results of phytochemical screening were also found most similar in the phytochemicals composition of different solvents (n-butanol, chloroform, acetone, ethanol, and methanol). Alkaloids, flavonoids, carbohydrates, and diterpenes were detected positive in n-butanol extract of non-diazotrophic grown *Nostoc muscorum*, because of less polarity of n-butanol. On the other side, phytochemical screening showed negative findings for glycosides, saponins, phenolics, and proteins. Alkaloids, carbohydrates, and diterpenes were present in the n-chloroform extract of non-diazotrophic grown *Nostoc muscorum*. Because of lower chloroform polarity, Phytochemical screening showed negative findings for glycosides, saponins, phenolics compound, and proteins. The results of phytochemical screening were also showed most similarity in the butanol and chloroform extract. Alkaloids, flavonoids, carbohydrates, and diterpenes were present while glycoside, saponins, phenolics, and proteins were absent in acetone extract. Similar results were found in methanolic and ethanolic extract as alkaloids, alkaloid flavonoids, carbohydrates, and diterpenes were present while glycoside, phenolics, and proteins were absent in acetone extract. Saponins were absent in the ethanolic extract of non-diazotrophically grown *Nostoc muscorum*. Saponins were absent in the ethanolic extract of non-diazotrophically grown *Nostoc muscorum*.

S. No.	Constituents	N-butanol extract	Chloroform Extract	Acetone Extract	Methanol extract	Ethanol extract
1.	Alkaloids Hager's Test:	+	+	+	+	+
2.	Glycosides Legal's Test:	-	-	-	-	-
3.	Flavonoids Lead acetate Test:	+	+	+	+	+
4.	Saponins Froth Test:	-	-	-	+	-
5.	Phenolics Ferric Chloride Test:	-	-	-	-	-
6.	Carbohydrate Fehling's Test:	+	+	+	+	+
7.	Proteins Xanthoproteic Test:	-	-	-	-	-
8.	Diterpenes Copper acetate Test:	+	+	+	+	+

Table no. 2: Result of Phytochemical Screening of NH₄Cl sample.

The result of the phytochemical screening of n-butanol and chloroform extract of diazotrophically grown *Nostoc muscorum* are depicted in table no. 3. The results of phytochemical screening showed similarity in the butanol and chloroform extract. Alkaloids, Flavonoids, carbohydrates, and diterpenes were found to be present while glycoside, saponins, phenolics, and proteins were absent in both the extract. Non-diazotrophically grown *Nostoc muscorum* were also showed similar results in acetone extract where alkaloids, flavonoids and carbohydrates were found to be present while glycoside, saponins, phenolics and protein, and diterpenes were absent in acetone extract. In the methanol extract, Alkaloids, flavonoids, carbohydrates, and diterpenes were present while glycoside, saponins, and protein were absent in acetone extract. In ethanol, Alkaloid, flavonoids, carbohydrates, and diterpenes were found to be present while glycoside, saponins, phenolic, and proteins were absent in acetone extract.

Sr. No.	Constitute	N-butanol extract	Chloroform extract	Acetone extract	Methanol extract	Ethanol extract
1	Alkaloids Hager's Test:	+	+	+	+	+
2.	Glycosides Legal's Test:	-	-	-	-	-
3.	Flavonoids Lead acetate Test:	+	+	+	+	+
4.	Saponins Froth Test:	-	-	-	+	-
5.	Phenolics Ferric Chloride Test:	-	-	-	-	-
6.	Carbohydrate Fehling's Test:	+	+	+	+	+
7.	Proteins Xanthoproteic Test:	-	-	-	-	-
8.	Diterpenes Copper acetate Test:	+	+	-	+	+


Table no. 3: Result of the phytochemical screening without NH₄Cl sample.

3.2 Thin layer chromatography

Thin layer chromatography profile of different extracts of n-butanol, chloroform, methanol, and ethanol non-diazotrophic extract with toluene: Ethyl acetate: Formic acid (5:4:1 v/v/v) as mobile phase resolved major spots at different R_f at 254nm (short wavelength) and normal light.

Sample extract	n-butanol, chloroform, acetone, methanol, and ethanol non-diazotrophs extract				
Mobile Phase	Toluene: Ethyl acetate: Formic acid (ratio= 5:4:1)				
Plate Tag	NH ₄ Cl				
Distance traveled by mobile phase	5.0 cm				
<u>Number of spots</u>					
	n-butanol	Chloroform	Acetone	Ethanol	Methanol
Normal Light	4	5	5	2	4
Short Wavelength	4	5	5	2	4
Long Wave Length	4	5	6	4	6
Visibility	Considerable				
Any Match with Quercetin Standard	Yes (0.60cm.)				
<u>R_f Values of each spot (from bottom to top)</u>					
	n-butanol	Chloroform	Acetone	Ethanol	Methanol
Normal Light	0.58,0.64,	0.6,0.66,0.4 0.90,0.98	0.6,0.66,0.7 2.0,0.92,0.98	0.6,0.66	0.6,0.66,0.8 8.0,0.92

Short Wavelength	0.58,0.64, 0.90,0.98	0.6,0.66,0.4 0.90,0.98	0.6,0.66,0.7 2,0.92,0.98	0.6,0.66	0.6,0.66,0.8 8,0.92
Long Wave Length	0.58,0.64, 0.90,0.98	0.6,0.66,0.4 0.90,0.98	0.6,0.66,0.7 2,0.84,0.92, 0.98	0.6,0.66,0.86,0 .94	0.6,0.64,0.7 6,0.88,0.92
Spot Sequence (Left to right)					
First	Quercetin				
Second	n-butanol extract				
Third	Chloroform extract				
Fourth	Acetone extract				
Fifth	Ethanol extract				
Sixth	Methanol extract				

Table no. 4: Results of thin layer chromatography of non-diazotrophic extracts.

Normal light


Short UV

Long UV

Figure 1: TLC of n-butanol, chloroform, acetone, methanol, and ethanol non-diazotrophs extract.

Sample extract	n-butanol, chloroform, acetone, methanol, and ethanol non-diazotrophs extract				
Mobile Phase	Toluene: Ethyl acetate: Formic acid (ratio= 5:4:1)				
Plate Tag	without NH ₄ Cl				
Distance traveled by mobile phase	5.0 cm				
Number of spots					
	n-butanol	Chloroform	Acetone	Ethanol	Methanol
Normal Light	2	5	5	4	6
Short Wavelength	2	5	4	2	6
Long Wave Length	5	5	6	5	8
Visibility	Considerable				
Any Match with Quercetin Standard	Yes (0.60cm.)				
R _f Values of each spot (from bottom to top)					
	n-butanol	Chloroform	Acetone	Ethanol	Methanol
Normal Light	0.6,0.76	0.6,0.66, 0.74,0.9, 0.98	0.58,0.62, 0.74,0.9, 0.98	0.58,0.64, 0.76,0.88	0.58,0.62, 0.72,0.82, 0.88,0.96
Short Wavelength	0.6,0.76	0.6,0.66, 0.74,0.9, 0.98	0.58,0.62, 0.74,0.9	0.58,0.64	0.58,0.62, 0.72,0.82, 0.88,0.96
Long Wave Length	0.6,0.66, 0.76,0.84, 0.9	0.6,0.66, 0.74,0.9, 0.98	0.58,0.62, 0.74,0.8, 0.9,0.98	0.58,0.64,0.86,0 .9, 0.98	0.58,0.62, 0.72,0.76, 0.82,0.88, 0.9,0.96
Spot Sequence (Left to right)					
First	Quercetin				
Second	n-butanol extract				
Third	Chloroform extract				
Fourth	Acetone extract				

Fifth	Ethanol extract
Sixth	methanol extract

Table no. 5: Results of thin layer Chromatography of diazotrophs extracts.**Figure 2:** TLC of n-butanol, chloroform, acetone, methanol, and ethanol diazotrophs extract.

4. CONCLUSION

Based on the above findings, it may be concluded that organic solvent serves as the most capable solvent when compare to others. Therefore, Cyanobacterial extract could be seen as an admirable source of pharmaceutical purpose. With this regard, we will be able to develop new drugs from *Nostoc muscorum* and this new drug would be cheap and more effective against pathogens. And thus it will help in removing mycobacterium tuberculosis from this Indian society.

ACKNOWLEDGMENT

SD is very thankful to the Head of the Department of Zoology for providing the necessary facilities. I am also thankful to the all staff members for their support and encouragement.

REFERENCES

- Biondi N, Piccardi R, Margheri MC, Rodolfi L, Smith GD, Tredici MR. Evaluation of *Nostoc* strain ATCC 53789 as a potential source of natural pesticides. *Applied and environmental microbiology*. 2004 Jun;70(6):3313-20.
- Dahms HU, Ying X, Pfeiffer C. Antifouling potential of cyanobacteria: a mini-review. *Biofouling*. 2006 Jan 1;22(5):317-27.
- Fish SA, Codd GA. Bioactive compound production by thermophilic and thermotolerant cyanobacteria (blue-green algae). *World Journal of Microbiology and Biotechnology*. 1994 May;10(3):338-41.
- Gerloff GC, Fitzgerald GP, Skoog F. The isolation, purification, and culture of blue-green algae. *American journal of botany*. 1950 Mar 1;216-8.
- Ghasemi Y, Yazdi MT, Shafiee A, Amini M, Shokravi S, Zarrini G. Parsiguine, a novel antimicrobial substance from *Fischerella ambigua*. *Pharmaceutical biology*. 2004 Jan 1;42(4-5):318-22.

- Noaman NH, Fattah A, Khaleafa M, Zaky SH. Factors affecting antimicrobial activity of *Synechococcus leopoliensis*. *Microbiological Research*. 2004 Dec 15;159(4):395-402.
- Metting B, Pyne JW. Biologically active compounds from microalgae. *Enzyme and Microbial Technology*. 1986 Jul 1;8(7):386-94.
- Pflugmacher S. Possible allelopathic effects of cyanotoxins, with reference to microcystin-LR, in aquatic ecosystems. *Environmental Toxicology: An International Journal*. 2002;17(4):407-13.
- Whitton BA, Potts M, editors. *The ecology of cyanobacteria: their diversity in time and space*. Springer Science & Business Media; 2007 May 8.
- Rodríguez-Ezpeleta N, Brinkmann H, Burey SC, Roure B, Burger G, Löffelhardt W, Bohnert HJ, Philippe H, Lang BF. Monophyly of primary photosynthetic eukaryotes: green plants, red algae, and glaucophytes. *Current biology*. 2005 Jul 26;15(14):1325-30.