

Environmental pollutants in organic farming and organic food**Kartikay Bisen, Udai Bhan Singh, Rahul Singh Rajput, Ankit Tiwari, Jitendra Kumar,
Raghvendra Singh, Aneeta Yadav**

Faculty of Agricultural Sciences and Allied Industries, Rama University, Kanpur-209217

Corresponding author email: kartikaybisen.fas@ramaunivrsity.ac.in**Abstract**

In current world scenario organic farming is a famous agricultural practice and covering a total area of approximately 71.5 Mha worldwide. Due to the involvement of certain practices in organic farming, food safety and human health is threatened by the flows of various environmental pollutants. Persistent organic pollutants, heavy metals, cyclodienes, organochlorine pesticides, hexachlorocyclohexanes, hexachlorobenzene, and nonbrominated flame retardants are some of the contaminants that occur in higher concentration in organic products. As a result, the lasting consumptions of organic products carrying such pollutants cause a considerable risk to human health. This present article decisively deal with the effect of organic farming on unknown and hidden threat due to the use of various composts as the nutrient and amendment sources that augment the phytoaccumulation and trophic transfer of pollutants, the functional diversity of the ecosystems, and poor harmonization among the policies and regulations in different countries for organic farming.

1. Introduction

Organic farming is widely practiced form of agriculture and suffers from poor attentiveness about the many possible means of entry and flow of pollutants. The flame-retardants (FRs), heavy metals, veterinary antibiotics (VAs), and even antimicrobial-resistant microorganisms are the new threats due to existing organic farming practices (Manyi-Loh et al., 2018). Many antibiotic resistant animal bacteria can be pathogens and furthermore the distribution of these bacteria from animal excretes to the soil and local water body can easily move to the human via the food chain (Manyi-Loh et al., 2018). Various organic compounds and pesticides residues as diffuse pollutants have been found in organic farm produce. The HPLC MS analysis detected many pesticides in organic carrots (Chiarello and Moura, 2018).

This inadvertent flow of pollutants into organic produce can be credited to using a range of farm input like farmyard manure, green waste, compost, domestic waste, sludge, and forage grass at organic farm. These manures are the main sources of VAs, pesticides and their residues, pharmaceutical compounds, microplastics, personal care products, and heavy metals (Umlauf et al., 2011; Meng et al., 2017; Zhang et al., 2020). The sustainable practices of organic farming have expanded enough, while human health benefits require the careful consideration of conventional and emerging pollutants. Knowledge on the probable flow of pollutants in organic farming is vital to develop a system and policy regulations to guaranteed food security and human health.

2. Composting: unintended piling of pollutants

Farm animal waste based manures that may contain steroid hormone metabolites have shown noteworthy unfavorable effects on the soil (Michelini et al., 2014; Verderame et al., 2016). The uses of antibiotics in farm animals are common worldwide (Tasho and Cho, 2016; Quaik et al., 2020).

Later, the direct livestock excretion on the land and the application of farm manure in agriculture, coupled with surface runoff and infiltration into deep layers of soils or water can release VAs into the soil and water system (Kemper, 2008; Kuppusamy et al., 2018). These reports suggest that the application of farm animal manure in organic farming poses unseen hazard and more importantly, the entry of VAs into the food chain. These threats significantly necessitate the screening of farm yard manures before its application in agricultural fields.

Table 1 Environmental pollutants in the agricultural inputs used as nutrient

Input	Pollutants	Reference
Swine manure	\sum_{15} VAs	Zhi et al. (2020)
Dairy manure	\sum TCs	Li et al. (2020)
Beef manure	\sum TCs	Li et al. (2020)
Chicken manure	\sum TCs	Li et al. (2020)
Sludge	Heavy metals; Cu, Zn, Cr, Pb, Mn, Ni	Tang et al. (2020)

Dried sludge	Heavy metals; Cu, Zn, Cr, Pb, Mn, Ni	Chen et al. (2019)
Tap water	∑32PPCPs	Liu et al. (2019)
Urban river water	Sulpiride	Tamura et al. (2017)
Dairy cattle manure	Ciprofloxacin	Gros et al. (2019)

Green waste or biological waste is usually consisting of farm and agriculture refuse (e.g., grass clippings and leaves), domestic or industrial kitchen waste. Farm waste is commonly used in organic farming owing to its high nitrogen content, and mitigating potential of greenhouse gas emissions (Diacono et al., 2019). Nevertheless, green wastes that have persistent herbicides residues pose health risks to organic farm produces (CalRecycle, 2020). The pesticides residues have been detected in various foods and the maximum residual level in organic food stuffs were reported more than 0.5% (Tiryaki, 2017). In recent times, the certified organic grains produced in the USA were reported to be contaminated with herbicide glyphosate (Meftaul et al., 2020). In general, use of synthetic pesticides are restricted in organic farming, but however the organic certified produce might contain the trace amount of pesticides which may be about five times lower than the amount detected in the conventional produce (Gómez-Ramos et al., 2020).

Urban sewage sludge is used as an organic amendment in organic farming and which provide organic matter and N and P. The presence of high organic matter in the sludge absorbs heavy metals from sewage during the treatment process. Thus, the hidden danger with the use of urban sewage sludge in organic farming is the presence of heavy metals (Table-1). The heavy metals can readily enter the plant system from this type of organic amendments, and their accumulation in plants can exceed the permissible levels for human consumption (Weldegebriel et al., 2012).

Table 2 Heavy metals in foodstuffs

Bovine meat	DL-PCBs	Tressou et al. (2017)
Bovine, poultry and lamb meat	PAHs, PCBs, OCPs	Hernández et al. (2017)

Cold-pressed rapeseed oil	PCBs, PAHs, Pesticides, Trace elements	Wroniak and Rekas (2017)
Vegetables	Cd, Co, Cr, Cu, Fe, Mn, Ni, Pb, Zn, Micronutrients	Hattab et al. (2019)
Carrots	Heavy metals	Gawęda et al. (2010)
Cereals, flours and bakery products	Ochratoxin A	Beretta et al. (2002); Biffi et al. (2004)
Cereals and cereal-based products	Mycotoxins	Pleadin et al. (2017)
Eggs	PCBs, PCDD/Fs	Rawn et al. (2012)
Cheese	PCBs, OCPs	Almeida-González et al. (2012)
Infant cereals	Cd	Hernández-Martínez and Navarro-Blasco (2012)

3. Knowledge gaps and future prospects

Knowledge about the various classes of heavy metals in the organic wastes is inadequately presented, limited to a very few research (Kolesnikov et al., 2019). The heavy metals in present different forms may have distinct levels of toxicities (Kolesnikov et al., 2019). In many cases certain pollutants and contaminants below the legal limits are found to be toxic to the plant and human. The extent of heavy metals pollutants in the organic sludge wastes alone can't be associated to toxicities. Regulatory guidelines for eco-toxicity by heavy metals are very tricky to define because of this limitation. Thus, future studies must focus on the speciation of heavy metals pollutants in the organic wastes and manure used in organic farming.

There should be an additional spotlight on the plant uptake of antibiotics from organic wastes. The research with antibiotics in hydroponic system cannot offer complete information about the plant uptake. The inconsistency and doubt between the translocation factors and antibiotic hydrophobicity need more research attention (Miller et al., 2016).

Organic wastes as soil amendment as a fertilizer in organic farming is usually considered secure. Nevertheless, techniques for the detection of pollutants are careful monitoring. The currently available treatment methods may not remove certain environmental pollutants like PPCPs. The emerging analytical techniques such as photoelectrochemical, complex nanocomposites, 3D flower-like ZnO nanoparticles (NPs), aptamer based colorimetric sensing, rhodamine assisted fluorescent approach, Cd-MOF, surface plasmon resonance (SPR) sensor, microbial fuel cell (MFC) biosensors may improve the detection and monitoring of pollutants.

4. Conclusion

The unintended stream of environmental pollutants can take place through compost and different other organic sources. The trophic transport of pollutants has many serious consequences in the agriculture production ecosystem. The environmental pollutants including heavy metals, pesticides, PPCPs in the organic compost, green waste, urban and sewage waste can reach to the agricultural fields. The organic plant products were found to have ultra-trace levels of pesticides, antibiotics, and heavy metals. Incompetent treatment methods of sludge and poor treatment of organic wastes can result in the flow and continuous existence of environmental pollutants in the organic farming fields. The validation of organic farming for enhancing the biodiversity becomes difficult to encourage the producers and policymakers. The further research and policy regulations are required to guarantee the unintended flow of environmental pollutants into organic products and keep the health of humans and ecosystems.

Reference

Almeida-González, M., Luzardo, O.P., Zumbado, M., Rodríguez-Hernández, Á., Ruiz-Suárez, N., Sangil, M., Camacho, M., Henríquez-Hernández, L.A., Boada, L.D., 2012. Levels of organochlorine contaminants in organic and conventional cheeses and their impact on the health of consumers: an independent study in the Canary Islands (Spain). *Food Chem.Toxicol.* 50, 4325–4332.

Beretta, B., Domenico, R.D., Gaiaschi, A., Ballabio, C., Galli, C.L., Gigliotti, C., Restani, P., 2002. Ochratoxin a in cereal-based baby foods: occurrence and safety evaluation. *Food Addit. Contam.* 19, 70–75.

Biffi, R., Munari, M., Dioguardi, L., Ballabio, C., Cattaneo, A., Galli, C.L., Restani, P., 2004. Ochratoxin A in conventional and organic cereal derivatives: a survey of the Italian market, 2001–02. *Food Addit. Contam.* 21, 586–591.

CalRecycle. 2020. Pesticide/herbicide residues in compost. <https://www.calrecycle.ca.gov/organics/threats/pesticides> (accessed 14 October 2020).

Chen, L., Liao, Y., Ma, X., Lu, S., 2020. Heavy metals chemical speciation and environmental risk of bottom slag during co-combustion of municipal solid waste and sewage sludge. *J. Clean. Prod.* 262, e121318.

Chiarello, M., Moura, S., 2018. Determination of pesticides in organic carrots by highperformance liquid chromatography/high-resolution mass spectrometry. *Anal. Lett.* 51, 2563–2576.

Diacono, M., Persiani, A., Testani, E., Montemurro, F., Ciaccia, C., 2019. Recycling agricultural wastes and by-products in organic farming: biofertilizer production, yield performance and carbon footprint analysis. *Sustainability* 11, e3824.

Gawęda, M., Nizioł-Łukaszewska, Z., Szopińska, A., 2010. The contents of selected metals in carrot cultivated using conventional, integrated and organic method. *Acta Hortic.* 936, 257–263.

Gómez-Ramos, Md.M., Nannou, C., Bueno, M.J.M., Goday, A., Murcia-Morales, M., Ferrer, C., Fernández-Alba, A.R., 2020. Pesticide residues evaluation of organic crops. A critical appraisal. *Food Chem.* 5, e100079.

Gros, M., Mas-Pla, J., Boy-Roura, M., Geli, I., Domingo, F., Petrović, M., 2019. Veterinary pharmaceuticals and antibiotics in manure and slurry and their fate in amended agricultural soils: findings from an experimental field site (Baix Empordà, NE Catalonia). *Sci. Total Environ.* 654, 1337–1349.

Hattab, S., Bougattass, I., Hassine, R., Dridi-Al-Mohandes, B., 2019. Metals and micronutrients in some edible crops and their cultivation soils in eastern-central region of Tunisia: a comparison between organic and conventional farming. *Food Chem.* 270, 293–298.

Hernández, Á.R., Boada, L.D., Mendoza, Z., Ruiz-Suárez, N., Valerón, P.F., Camacho, M., Zumbado, M., Almeida-González, M., Henríquez-Hernández, L.A., Luzardo, O.P., 2017. Consumption of organic meat does not diminish the carcinogenic potential associated with the intake of persistent organic pollutants (POPs). *Environ. Sci. Pollut. Res.* 24, 4261–4273.

Hernández-Martínez, R., Navarro-Blasco, I., 2012. Estimation of dietary intake and content of lead and cadmium in infant cereals marketed in Spain. *Food Control* 26, 6–14.

- Kemper, N., 2008. Veterinary antibiotics in the aquatic and terrestrial environment. *Ecol. Indic.* 8, 1–13.
- Kolesnikov, S.I., Timoshenko, A.N., Kazeev, K.S., Akimenko, Y.V., Soldatov, A.V., 2019. Comparison of ecotoxicity of nickel and iron oxides and their nanoforms. *Rasayan J. Chem.* 12, 549–553.
- Kuppusamy, S., Kakarla, D., Venkateswarlu, K., Megharaj, M., Yoon, Y.-E., Lee, Y.B., 2018. Veterinary antibiotics (VAs) contamination as a global agro-ecological issue: a critical view. *Agric. Ecosyst. Environ.* 257, 47–59.
- Li, C., Li, Y., Li, X., Ma, X., Ru, S., Qiu, T., Lu, A., 2020. Veterinary antibiotics and estrogen hormones in manures from concentrated animal feedlots and their potential ecological risks. *Environ. Res.* 13, e110463.
- Liu, M., Yin, H., Wu, Q., 2019. Occurrence and health risk assessment of pharmaceutical and personal care products (PPCPs) in tap water of Shanghai. *Ecotoxicol. Environ. Saf.* 183, e109497.
- Manyi-Loh, C., Mamphweli, S., Meyer, E., Okoh, A., 2018. Antibiotic use in agriculture and its consequential resistance in environmental sources: potential public health implications. *Molecules (Basel, Switzerland)*. 23, e795.
- Meftaul, I.Md., Venkateswarlu, K., Dharmarajan, R., Annamalai, P., Asaduzzaman, Md, Parven, A., Megharaj, M., 2020. Controversies over human health and ecological impacts of glyphosate: is it to be banned in modern agriculture? *Environ. Pollut.* 263, 114372.
- Meng, J., Wang, L., Zhong, L., Liu, X., Brookes, P.C., Xu, J., Chen, H., 2017. Contrasting effects of composting and pyrolysis on bioavailability and speciation of Cu and Zn in pig manure. *Chemosphere.* 180, 93–99.
- Michelini, L., Gallina, G., Capolongo, F., Ghisi, R., 2014. Accumulation and response of willow plants exposed to environmental relevant sulfonamide concentrations. *Int. J. Phytoremediation* 16, 947–961.
- Miller, E.L., Nason, S.L., Karthikeyan, K.G., Pedersen, J.A., 2016. Root uptake of pharmaceuticals and personal care product ingredients. *Environ. Sci. Technol.* 50, 525–541.
- Pleadin, J., Staver, M.M., Markov, K., Frece, J., Zadravec, M., Jaki, V., Krupić, I., Vahčić, N., 2017. Mycotoxins in organic and conventional cereals and cereal products grown and marketed in Croatia. *Mycotoxic Res.* 33, 219–227.

- Quaik, S., Embrandiri, A., Ravindran, B., Hossain, K., Al-Dhabi, N.A., Arasu, M.V., Ignacimuthu, S., Ismail, N., 2020. Veterinary antibiotics in animal manure and manure laden soil: scenario and challenges in Asian countries. *J. King Saud. Uni. Sci.* 32, 1300–1305
- Rawn, D.F., Sadler, A.R., Quade, S.C., Sun, W.F., Kosarac, I., Hayward, S., Ryan, J.J., 2012. The impact of production type and region on polychlorinated biphenyl (PCB), polychlorinated dibenzo-p-dioxin and dibenzofuran (PCDD/F) concentrations in Canadian chicken egg yolks. *Chemosphere* 89, 929–935.
- Tamura, I., Yasuda, Y., Kagota, K.I., Yoneda, S., Nakada, N., Kumar, V., Kameda, Y., Kimura, K., Tatarazako, N., Yamamoto, H., 2017. Contribution of pharmaceuticals and personal care products (PPCPs) to whole toxicity of water samples collected in effluentdominated urban streams. *Ecotoxicol. Environ. Saf.* 144, 338–350.
- Tang, J., He, J., Tang, H., Wang, H., Sima, W., Liang, C., Qiu, Z., 2020. Heavy metal removal effectiveness, flow direction and speciation variations in the sludge during the biosurfactant-enhanced electrokinetic remediation. *Sep. Purif. Technol.* 246, e116918.
- Tasho, R.P., Cho, J.Y., 2016. Veterinary antibiotics in animal waste, its distribution in soil and uptake by plants: a review. *Sci. Total Environ.* 563-564, 366–376.
- Tiryaki, O., 2017. Pesticide residues and organic production. *J. Biol. Environ. Sci.* 11, 11–23
- Tressou, J., Abdallah, N.B., Planche, C., Dervilly-Pinel, G., Sans, P., Engel, E., Albert, I., 2017. Exposure assessment for dioxin-like PCBs intake from organic and conventional meat integrating cooking and digestion effects. *Food Chem. Toxicol.* 110, 251–261.
- Umlauf, G., Christoph, E.H., Lanzini, L., Savolainen, R., Skejo, H., Bidoglio, G., Clemens, J., Goldbach, H., Scherer, H., 2011. PCDD/F and dioxin-like PCB profiles in soils amended with sewage sludge, compost, farmyard manure, and mineral fertilizer since 1962. *Environ. Sci. Pollut. Res.* 18, 461–470
- Verderame, M., Limatola, E., Scudiero, R., 2016. Estrogenic contamination by manure fertilizer in organic farming: a case study with the lizard *Podarcis sicula*. *Ecotoxicology* 25, 105–114.
- Weldegebriel, Y., Chandravanshi, B.S., Wondimu, T., 2012. Concentration levels of metals in vegetables grown in soils irrigated with river water in Addis Ababa. Ethiopia. *Ecotoxicol. Environ. Saf.* 77, 57–63.
- Wroniak, M., Rekas, A., 2017. A preliminary study of PCBs, PAHs, pesticides and trace metals contamination in cold-pressed rapeseed oils from conventional and ecological cultivations. *J. Food Sci. Technol.* 54, 1350–1356.

Zhang, L., Xie, Y., Liu, J., Zhong, S., Qian, Y., Gao, P., 2020. An overlooked entry pathway of microplastics into agricultural soils from application of sludge-based fertilizers. *Environ. Sci. Technol.* 54, 4248–4255.

Zhi, S., Zhou, J., Liu, H., Wu, H., Zhang, Z., Ding, Y., Zhang, K., 2020. Simultaneous extraction and determination of 45 veterinary antibiotics in swine manure by liquid chromatography-tandem mass spectrometry. *J. Chromatogr. B* 1154, e122286.