



Volume 8, Issue 2,

April 2019,

www.ijfans.com

e-ISSN: 2320-7876

**INTERNATIONAL JOURNAL OF FOOD AND
NUTRITIONAL SCIENCES**

IMPACT FACTOR ~ 1.021



Official Journal of IIFANS

EFFECTS OF CHELATING AGENTS ON LIPID OXIDATION, PH, AND COLOR CHANGE IN REDUCED SODIUM AND LOW-FAT PORK PATTIES

Shu-Tai Wang¹ and Jen-Hua Cheng^{2*}

*Corresponding Author: **Jen-Hua Cheng**, ✉ jhcheng115@yahoo.com.tw

Received on: 19th January, 2019

Accepted on: 11th March, 2019

The purpose of this study was to evaluate the effectiveness of adding chelating agents to inhibit lipid oxidation and color change in precooked pork patties. The reduced sodium and low fat patties were used because of health concern by consumers. The pork patties with EDTA had a significantly highest Hunter b value compared to the other samples. The Hunter L values of precooked pork patties were significantly higher at day 7 when compared to that at day 0. The redness of precooked pork patties decreased during refrigerated storage, except for the sample with 0.5% sodium tripolyphosphate. There was a significant interaction between treatment and storage time for TBA values. During refrigerated storage, TBA values of samples with chelating agents maintained lipid oxidation up to 7 days. According to these results, addition of chelating agents did solve the problems of lipid oxidation and color evaluation in reduced sodium and low-fat precooked pork samples.

Keywords: Chelating agents, Lipid oxidation, Color change, Pork patties

INTRODUCTION

Today, consumers not only care about flavor and convenience of food, but are also concerned with hygiene, safety, and nutrition of food. The consumption of food has changed and healthy food is popular. The consumers like to eat low-fat food (Egbert *et al.*, 1991). However, fat increases the juiciness and tenderness of meat, and it affects the flavor, mouthfeel, and color of meat.

Sodium chloride is usually used in the meat curing process. Its functions include inhibition of microorganisms, increase of flavor, and extraction of salt soluble proteins. Usually, sodium chloride is added at 2.5-3.0% of cured product weight in meat products. However, the sodium ion can cause hypertension; therefore, it has been recommended to reduce daily intake of salt. Thus, a new method has been developed to reduce the amount of sodium in meat. The

reduction of sodium chloride in meat products causes loss of cooking yield, changes in flavor and shorter shelf life (Sofos, 1986).

Lipid oxidation causes the quality of meat to change, produces off-flavor, and decreases flavor, color, texture, nutritive value, and acceptance (Buckley *et al.*, 1989). Autoxidation is one aspect of lipid oxidation, which is a chain reaction of free radicals, and can be separated into three steps: initiation, propagation and termination (Min, 1998). The autoxidation of unsaturated fatty acids produce potentially toxic compounds (Addis and Park, 1989).

Chelating agents have been used to tie up metals in the initiation of autoxidation. Chelating agents bind with heavy metals in the initiation of autoxidation and cause metal ions to lose their ability to catalyze lipid oxidation. It is an effective method to inhibit lipid oxidation by attacking metal ions

¹ Department of Hospitality Management, Tunghai University, Taichung, Taiwan.


² Department of Nutrition and Health Sciences, Kainan University, Taoyuan City 338, Taiwan.

directly (Gorji *et al.*, 2016). The metal ions in muscle, especially iron ions, catalyze lipid oxidation (Min, 1998). The basic hypothesis of this study is the utilization of chelating agents which bind with metal ions, and it is a useful method to inhibit lipid oxidation. There is little study about chelating agents using in reduced sodium and low-fat meat system.

The purpose of this study was to evaluate the effectiveness of adding chelating agents for inhibiting lipid oxidation and color changes in reduced salted and low-fat pork patties.

MATERIALS AND METHODS

The 24 hours post-mortem pork ham was ground through a 3 mm plate and then divided into 3 groups. NaCl was added at the level of 0.5%, and the raw meat was previously blended with chelating agents (no addition, 0.5% sodium tripolyphosphate or 500 ppm EDTA). Pork patties were prepared by molding 90 gm of ground pork in a Petri dish with a diameter of 90 mm and the thickness of patties were 5 mm. Pork patties were cooked on both sides for a total of 20 minutes on a hot plate until the internal temperature of the patties reached 75 °C. These precooked pork patties were placed in a tray, covered with aluminum foil, and stored at 4 °C. Measurements of cooking yield, TBARS values, a*, b*, L* color values and pH value were tested on precooked pork patties at 0 and 7 days of storage. This test was repeated 3 times (Figure 1).



Cooking Yield

Cooking Yield = (final product weight/fresh meat weight)*100%

pH Measurement

The pH values of samples were measured with a pH meter (Ockerman, 1985). The pH meter was standardized by buffer solutions (pH 7.00 and 4.00 at 25 °C) before testing. Ten grams of sample was weighed and blended with 100 ml of distilled water in a polyethylene sterile bag by a Stomacher (Easy Mix, Germany) for 1 minute.

TBARS Values

The sample was analyzed by the modified extraction method of TBARS (Pensel, 1990). A 5-gram sample of meat tissue was placed in a polyethylene bag. Fifty ml of chilled (4 °C) solution of 20% trichloroacetic acid in 1.6% of phosphoric acid was added to the same bag and then massaged for two minutes in a Stomacher to mix the sample. Fifty ml of chilled distilled water (4 °C) was then added into the bag and the stomacher was again used to blend the sample for 30 seconds. The slurry was filtered through Whatman No. 1 filter paper into a 100 ml cylinder. Five ml of the filtrate was pipetted into a test tube, and five ml of fresh chilled 0.02 M 2-thiobarbituric acid solution was added to this tube. All samples in test tubes were placed in the dark at room temperature (25 °C) for 15 hours in order to develop the color reaction. The amount of color was measured using a UV/VIS spectrophotometer (SP8001, Metertech Inc., Taiwan) at 532 nm to calculate the TBARS value.

Standard Curve of TBARS Test

Two point two grams of 1,1,3,3-tetraethoxypropane (TEP, FW = 220) was diluted to 1000 ml with distilled water to prepare a 10⁻² stock solution. A 10⁻³M stock solution of TEP was prepared by diluting on milliliter of the 10⁻² M TEP solution with nine milliliter of distilled water. Zero, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 and 0.9 ml volumes of the 10⁻³ (10⁻⁶

Table 1: Main Effect of b*, L* Values, pH Values of Reduced Sodium and Low-Fat Pork Patties with Chelating Agents During Refrigerated (4 °C) Storage

Main Effect Treatment	pH	b Value	L Value
	T1	6.44 ^{AB}	17.45 ^B
T2	6.55 ^A	19.13 ^B	40.79 ^B
T3	6.39 ^B	22.29 ^A	46.05 ^A
Storage Time			
Day 0	6.48 ^a	18.57 ^a	38.40 ^b
Day 7	6.43 ^a	20.68 ^b	43.54 ^a

Note: T1: Control, with no antioxidant; T2: with 0.5% sodium tripolyphosphate; T3: with 500ppm EDTA. ^{AB} Means with different uppercase superscripts within a column, within main effect of the treatments are significantly different (p<0.05). ^{ab} Means with different lowercase superscripts within a column, within main effect of storage time are significantly different (p<0.05).

moles of TEP) stock solution were pipetted into 50-ml volumetric flasks, and then diluted to volume with distilled water. Five ml of each diluted solution was mixed with 5 ml of 0.02 M TBA reagent in each test tube and placed in the dark to develop color for 15 hours. The absorbance of samples was measured by a spectrophotometer at 532 nm.

Color Measurement

Minolta colorimeter (CR-10, Konica Minolta Sensing, Inc., Japan) was used to detect a*, b* and L* values of samples. The a* measures redness; b* indicates yellowness; and L* indicates lightness. Each value was the mean of 10 determinations.

Statistical Analysis

Data were analyzed using the General Linear Model procedure of SAS. Differences among means were detected at the 5% level using Duncan's New Multiple Range Test by SAS.

RESULTS AND DISCUSSION

The effect of adding different antioxidants on the cooking yields of pork patties shows that there was no significant difference among the three treatments. Sodium tripolyphosphate and EDTA did not increase cooking yield compared to the control that had no added chelating agents.

There were no significant interactions between treatment and storage time for pH, b and L values; therefore, main effects are shown as results of these measurements. The addition of 0.5% sodium tripolyphosphate had higher ($p < 0.05$) pH value than samples with 500 ppm EDTA, but the pH values of pork patties with chelating agents were the same as control.

Treatment with sodium tripolyphosphate had the same Hunter b value compared to control ($P > 0.05$). The pork patties with EDTA had the significantly highest Hunter b value among samples. Also, the yellowness of precooked pork patties did increase during refrigerated storage ($p > 0.05$).

The control had significantly lower Hunter L value compared to samples with sodium tripolyphosphate that had the lowest yellowness value when compared to pork patties with EDTA ($p < 0.05$). The Hunter L values of precooked pork patties were significantly higher at day 7 compared to that at day 0.

There was significant interaction between treatment and storage time for Hunter a value. Treatments had the same

Hunter a values at day 0 and 7 ($p > 0.05$). However, the redness of precooked pork patties was significantly decreased during refrigerated storage, except for samples with 0.5% sodium tripolyphosphate (Table 2). Meat color is the major factor of acceptability by consumers.

There was significant interaction between treatment and storage time for TBA values. Table 3 shows that TBA values of precooked pork patties were affected by adding chelating

Table 2: Effect of Adding Chelating Agents on Hunter a* Values of Reduced Sodium and Low-Fat Pork Patties During Refrigerated (4 °C) Storage

Treatment	Storage Time	
	Day 0	Day 7
Control, 0%	13.06 ^{Aa}	10.73 ^{Ab}
	(0.46) ¹	(1.54)
0.5% Phosphate	13.14 ^{Aa}	11.97 ^{Aa}
	(1.65)	(2.68)
500 ppm EDTA	15.13 ^{Aa}	9.83 ^{Ab}
	(2.09)	(0.69)

Note: ¹ indicates standard deviation. ^A Different uppercase superscripts in the same column indicate significantly different ($p < 0.05$). ^{ab} Different lowercase superscripts in the same row indicate significantly different ($p < 0.05$).

Table 3: Effect of Adding Chelating Agents on TBARS1 Values of Reduced Sodium and Low-Fat Pork Patties During Refrigerated (4 °C) Storage

Treatment	Storage Time	
	Day 0	Day 7
Control, 0%	0.435 ^{Ab}	2.090 ^{Aa}
	(0.088) ²	(0.890)
0.5% Phosphate	0.303 ^{Ba}	0.360 ^{Ba}
	(0.048)	(0.127)
500 ppm EDTA	0.230 ^{Ba}	0.460 ^{Ba}
	(0.056)	(0.310)

Note: ¹ TBARS = mg malonaldehyde/kg muscle; ² indicates standard deviation. ^{AB} Different uppercase superscripts in the same column indicate significantly different ($p < 0.05$). ^{ab} Different lowercase superscripts in the same row indicate significantly different ($p < 0.05$).

agents. At day 0, samples with 500 ppm EDTA and with sodium tripolyphosphate (0.5%) had the significantly lower TBA values than the control; however, the TBA values of samples with chelating agents were at the same levels ($p > 0.05$). At day 7, control without any chelating agents had significantly higher TBA values than pork with sodium tripolyphosphate and EDTA. The TBA values of samples with sodium tripolyphosphate were the same as those of the EDTA treatment ($p > 0.05$). It seems that chelating agents significantly decreased lipid oxidation in precooked pork patties. Shahidi *et al.* (1985) reported that EDTA and sodium tripolyphosphate decreased TBA values in cooked meat. EDTA decreased lipid oxidation during refrigerated storage, presumably by its demonstrated effect on non-heme iron catalysis (Kwoh, 1971). The ability of chelating agent is to tie up metal ions in food system. The addition of EDTA, a well-known metal chelator, increased the oxidative stability in O/W emulsions (Yi *et al.*, 2016).

During refrigerated storage, TBA values of control were increased ($p < 0.05$); however, chelating agents maintained the lipid oxidation up to 7 days of refrigerated storage ($p > 0.05$). According to this result, the addition of 0.5% sodium tripolyphosphate and 500ppm EDTA effectively inhibited lipid oxidation in precooked pork patties. Adjimani and Asare (2015) reported that all the iron chelators were observed to show significant activities from studied antioxidant assays.

CONCLUSION

Chelating agents have been used to tie up metals in the initiation of autooxidation. The reduced sodium and low fat patties was used for health concerns of consumers. The redness of precooked pork patties was decreased during refrigerated storage, except for samples with 0.5% sodium tripolyphosphate. For TBA values, chelating agents significantly decreased lipid oxidation in precooked pork patties. According to these results, addition of chelating agents did solve problems of lipid oxidation and color evaluation in reduced sodium and low fat meat models.

REFERENCES

- Addis P B and Park S W (1989), *In Food Toxicology: A Perspective on the Relative Risks*, in S L Taylor and R A Scanlan (Eds.), p. 297, Marcel Dekker, New York.
- Adjimani J P and Asare P (2015), "Antioxidant and Free Radical Scavenging Activity of Iron Chelators", *Toxicology Reports*, Vol. 2, pp. 721-728.
- Buckley D J, Gray J I, Ashgar A, Price J F, Crackel R L, Booren A M, Pearson A M and Miller E R (1989), "Effects of Dietary Antioxidants and Oxidized Oil on Membral Lipid Stability and Pork Product Quality", *J. Food Sci. Off. Publ. Inst. Food Technol.*, Vol. 54, No. 5, pp. 1193-1197.
- Egbert W R, Huffman D L, Chen C and Dylewski D P (1991), "Development of Low Fat Ground Beef", *Food Tech.*, Vol. 45, No. 6, pp. 64-73.
- Gorji S G, Smyth H E, Sharma M and Fitzgerald M (2016), "Lipid Oxidation in Mayonnaise and the Role of Natural Antioxidants: A Review", *Trends in Food Science & Technology*, Vol. 56, pp. 88-102.
- Kwoh T L (1971), "Catalysts of Lipid Peroxidation in Meats", *Journal of American Oil Chemists' Society*, Vol. 48, No. 10, pp. 550-555.
- Min D B (1998), "Food Lipid: A Course Note", *Food Science & Technology*, pp. 57-71, The Ohio State University, Columbus, OH.
- Ockerman H W (1985), "Quality Control of Post-Mortem Muscle Tissue", Dept. of Animal Sciences, The Ohio State University, Columbus, OH.
- Pense N A (1990), "Influence of Experimental Conditions on Porcine Muscle and its Effect on Oxidation", A Thesis, The Ohio State University, Columbus, OH.
- SAS (2009), *Statistical Analysis System Package*, SAS Institute Inc., Cary, USA.
- Shahidi F, Rubin L J, Diosady L L, Kassam N, Li S F and Wood D F (1986), "Effect of Sequestering Agents on Lipid Oxidation in Cooked Meats", *Food Chemistry*, Vol. 21, No. 2, pp. 145-152.
- Sofos J N (1986), "Use of Phosphates in Low-Sodium Meat Products", *Food Technology*, Vol. 40, No. 9, pp. 52-64.
- Yi B R, Kim M J and Lee J H (2016), "Effects of Emulsifier Charges on the Oxidative Stability in Oil-in-Water Emulsions Under Riboflavin Photosensitization", *Food Science and Biotechnology*, Vol. 25, No. 4, pp. 1003-1009.

